

NEW PROOF OF THE $T(1)$ THEOREM FOR TRIEBEL–LIZORKIN SPACES

TUOMAS HYTÖNEN AND CORNELIA KAISER

Abstract. We give a new proof of the $T(1)$ theorem for reflexive homogeneous Triebel–Lizorkin spaces $\dot{F}_p^{s,q}$, which uses neither maximal functions nor atomic decompositions. The substitute tool is an estimate on translations in the $L_p(\ell_q)$ spaces.

2000 Mathematics Subject Classification: Primary: 47G10; Secondary: 42B20, 47B38.

Key words and phrases: Singular integral, translation operator, vector-valued estimate, Littlewood–Paley decomposition.

1. INTRODUCTION

After the publication of the remarkable $T(1)$ theorem of G. David and J.-L. Journé [1] concerning general conditions of L_2 boundedness of singular integral operators

$$Tf(x) = \int_{\mathbb{R}^N} K(x, y)f(y)dy,$$

several authors ([2, 3, 4, 5, 6]) proved similar results on various different function spaces such as homogeneous Besov spaces $\dot{B}_p^{s,q}$ and homogeneous Triebel–Lizorkin spaces $\dot{F}_p^{s,q}$ (which include spaces $L_p = \dot{F}_p^{s,2}$ for $1 < p < \infty$). In a different direction of generalization, T. Figiel [7] proved an analogue of the $T(1)$ theorem for X -valued functions $f \in L_p(X)$, where X is a Banach space with a certain additional property (UMD) but the kernel K is still scalar-valued. Quite recently, L. Weis and one of the present authors [8] gave a new proof of Figiel’s $T(1)$ theorem and extended it to operator-valued kernels K . The two main analytic ingredients in [8] are the well known Littlewood–Paley decomposition and a rather less known, but quite powerful, square-function estimate, due to J. Bourgain [9], for translation operators $\tau_h : f \mapsto f(\cdot - h)$. It was subsequently observed by the second-named author [10, 11] that the approach of [8] is adaptable to obtain proofs of $T(1)$ theorems also for vector-valued homogeneous Besov spaces $\dot{B}_p^{s,q}(X)$ and Bessel potential spaces $\dot{H}_p^s(X)$. The latter ones, in the scalar-valued case, coincide with $\dot{F}_p^{s,2}$ for $1 < p < \infty$.

The purpose of the present note is to show that essentially the same approach carries over to the whole scale of (reflexive) Triebel–Lizorkin spaces $\dot{F}_p^{s,q}$, $1 < p, q < \infty$. Thus, our aim is to give a new proof of the $T(1)$ theorem for these spaces which makes no use whatsoever of either maximal functions or

atomic decompositions, the methods employed in the earlier proofs of analogous results. The tool that replaces them in our proof is an appropriate modification to the present situation of the above-mentioned square function estimate for translations (see Lemma 3.1).

We conclude the introduction with our statement of the $T(1)$ theorem for the Triebel–Lizorkin spaces. The definitions of these spaces, as well as of various conditions appearing in the theorem, are given in Section 2. The crucial translation lemma is proved in Section 3, and the proof of Theorem 1.1 is then given in the last Section 4.

Theorem 1.1. *Let $n \in \mathbb{N}$ and $\nu \in (0, 1)$. Suppose $T \in \text{CZO}_{n+\nu}$ satisfies the weak boundedness property and the condition $T(u^\alpha) = 0$ for all $|\alpha| \leq n$. Then T extends to a bounded linear operator from $\dot{F}_p^{s,q}(\mathbb{R}^N)$ to $\dot{F}_p^{s,q}(\mathbb{R}^N)$ for each $s \in (0, n + \nu)$ and each $p, q \in (1, \infty)$.*

If, in addition, $T' \in \text{CZO}_{n+\nu}$ and $T'(u^\alpha) = 0$ for all $|\alpha| \leq n$, then the assertion holds for all $|s| < n + \nu$.

2. SPACES AND OPERATORS

We denote $\mathbb{N} := \{0, 1, 2, \dots\} \supseteq \mathbb{Z}_+ := \{1, 2, \dots\}$. We fix a number $N \in \mathbb{Z}_+$, and all our functions and distributions will be defined on \mathbb{R}^N . By $\mathcal{D}(\mathbb{R}^N) \subset \mathcal{S}(\mathbb{R}^N) \subset \mathcal{S}'(\mathbb{R}^N)$ we denote compactly supported smooth functions, rapidly decreasing smooth functions and the corresponding tempered distributions, respectively. The pairing of $\mathcal{S}(\mathbb{R}^N)$ and $\mathcal{S}'(\mathbb{R}^N)$ is denoted by $\langle \cdot, \cdot \rangle$. $\mathcal{Z}(\mathbb{R}^N)$ is the space of all Schwartz functions $\varphi \in \mathcal{S}(\mathbb{R}^N)$ such that $D^\alpha \widehat{\varphi}(0) = 0$ for all multiindices $\alpha \in \mathbb{N}^N$, where $\widehat{\varphi}$ is the Fourier transform of φ . Then $\mathcal{Z}(\mathbb{R}^N)$ is a closed subspace of $\mathcal{S}(\mathbb{R}^N)$. If $\mathcal{Z}'(\mathbb{R}^N)$ denotes the space of all continuous linear functionals on $\mathcal{Z}(\mathbb{R}^N)$, then $\mathcal{S}'(\mathbb{R}^N)/\mathcal{P}(\mathbb{R}^N)$ and $\mathcal{Z}'(\mathbb{R}^N)$ are isomorphic, where $\mathcal{P}(\mathbb{R}^N)$ is the space of polynomials in N real variables (cf. [12, 5.1.2]).

Homogeneous Triebel–Lizorkin spaces. Let $\widehat{\phi} \in \mathcal{D}(\mathbb{R}^N)$ be radial, equal to 1 in $\overline{B}(0, 1)$, and supported in $\overline{B}(0, 2)$. Let $\widehat{\varphi} = \widehat{\phi} - \widehat{\phi}(2\cdot)$ and $\widehat{\varphi}_{2^j} = \widehat{\varphi}(2^j\cdot)$, $j \in \mathbb{Z}$.

Let $p, q \in [1, \infty)$ and $s \in \mathbb{R}$. The *homogeneous Triebel–Lizorkin space* $\dot{F}_p^{s,q}(\mathbb{R}^N)$ is the space consisting of all $f \in \mathcal{Z}'(\mathbb{R}^N)$ such that

$$\|f\|_{\dot{F}_p^{s,q}(\mathbb{R}^N)} := \left\| \left(\sum_{j \in \mathbb{Z}} (2^{-js} |f * \varphi_{2^j}|)^q \right)^{1/q} \right\|_{L_p(\mathbb{R}^N)}$$

is finite. One can show that different choices of φ lead to equivalent norms and that $\dot{F}_p^{s,q}(\mathbb{R}^N)$, endowed with $\|\cdot\|_{\dot{F}_p^{s,q}(\mathbb{R}^N)}$, is a Banach space. Moreover $\mathcal{Z}(\mathbb{R}^N) \xrightarrow{d} \dot{F}_p^{s,q}(\mathbb{R}^N) \hookrightarrow \mathcal{Z}'(\mathbb{R}^N)$ (cf. [12, 5.1.5]).

Singular integral operators. Our main object of study is a continuous linear operator $T : \mathcal{S}(\mathbb{R}^N) \rightarrow \mathcal{S}'(\mathbb{R}^N)$. Its adjoint T' is a similar operator defined by $\langle \psi, T'\varphi \rangle := \langle \varphi, T\psi \rangle$.

Suppose that $K : \{(u, v) \in \mathbb{R}^N \times \mathbb{R}^N : u \neq v\} \rightarrow \mathbb{C}$ is a locally integrable function. We say that T is a *singular integral operator associated with K* if

$$\langle \varphi, T\phi \rangle = \int_{\mathbb{R}^N} \varphi(u) \int_{\mathbb{R}^N} K(u, v)\phi(v)dv du \tag{1}$$

holds for all $\varphi, \phi \in \mathcal{D}(\mathbb{R}^N)$ with $\text{supp}\varphi \cap \text{supp}\phi = \emptyset$. When this is the case, then T' is a singular integral operator with associated kernel K' given by $K'(u, v) = K(v, u)$ for $u \neq v$.

Next we introduce some more specific conditions on K and T :

The class $\text{CZO}_{n+\nu}$. Let $n \in \mathbb{N}$ and $\nu \in (0, 1)$. For a measurable kernel $K : \{(u, v) \in \mathbb{R}^N \times \mathbb{R}^N : u \neq v\} \rightarrow \mathbb{C}$, we consider the standard estimates

(SE _{n}) K is continuously differentiable up to order n with respect to the first variable and

$$\mathcal{C}_\alpha(K) = \sup\{|u - v|^{N+|\alpha|} |(\partial_u^\alpha K)(u, v)| : u \neq v\}$$

is finite for *all* multiindices $\alpha \in \mathbb{N}^N$ with $|\alpha| \leq n$;

(SE _{$n+\nu$}) K satisfies (SE _{n}) and

$$\mathcal{C}_{\alpha,\nu}(K) = \sup \left\{ |u - v|^{N+n+\nu} \frac{|(\partial_u^\alpha K)(u, v) - (\partial_u^\alpha K)(u_0, v)|}{|u - u_0|^\nu} : \begin{array}{l} |u - v| > 2|u - u_0| > 0 \end{array} \right\}$$

is finite for *some* multiindex $\alpha \in \mathbb{N}^N$ with $|\alpha| = n$.

We say that $T \in \text{CZO}_{n+\nu}$ if T is a singular integral operator associated with a kernel K satisfying (SE _{$n+\nu$}).

Note that $T \in \text{CZO}_{n+\nu}$ does *not* imply that $T' \in \text{CZO}_{n+\nu}$.

Definition of $T(u^\alpha)$. Let α be a multiindex with $|\alpha| \leq n$ and u^α the associated monomial. For $T \in \text{CZO}_{n+\nu}$, it can be shown (see [10]) that $u^\alpha T'\varphi$ agrees with an integrable function in the exterior of any neighbourhood of $\text{supp}\varphi$ provided that

$$\varphi \in \mathcal{D}^n(\mathbb{R}^N) := \{\varphi \in \mathcal{D}(\mathbb{R}^N) : \int_{\mathbb{R}^N} u^\beta \varphi(u)du = 0 \text{ for all } |\beta| \leq n\}.$$

Then we define

$$\langle \varphi, T(u^\alpha) \rangle := \langle \chi, u^\alpha T'\varphi \rangle + \langle 1 - \chi, u^\alpha T'\varphi \rangle,$$

where $\chi \in \mathcal{D}(\mathbb{R}^N)$ is any test function equal to unity in a neighbourhood of $\text{supp}\varphi$. The first pairing above is the usual one between a test function and a distribution, while the second one can be evaluated as a convergent Lebesgue integral. The definition is independent of χ and produces a well-defined object $T(u^\alpha) \in (\mathcal{D}^n(\mathbb{R}^N))'$.

The weak boundedness property. We say that φ is a *normalized bump function* associated with the unit ball if $\varphi \in \mathcal{D}(\mathbb{R}^N)$ with $\text{supp}\varphi \subseteq \overline{B}(0, 1)$ and $\|D^\alpha\varphi\|_\infty \leq 1$ for all $|\alpha| \leq M$, where M is a large fixed number. ϕ is a normalized bump function associated with the ball $\overline{B}(u, r)$ if $\phi(\cdot) = r^{-N}\varphi(r^{-1}(\cdot - u))$, where φ is a normalized bump function associated with the unit ball. The operator T has the *weak boundedness property* if for every pair of normalized bump functions φ, ϕ associated with any ball $\overline{B}(u, r)$ we have $|\langle \phi, T\varphi \rangle| \leq Cr^{-N}$.

The various conditions defined above will be used through the following implied estimates proved in [10] (see also [8]).

Lemma 2.1. *Let $k \in \mathbb{N}$, $a > 0$, $w \in \mathbb{R}^N$, and let $\varphi, \phi \in \mathcal{D}^0(\mathbb{R}^N)$ be normalized bump functions associated with $\overline{B}(0, a)$ and $\overline{B}(w, 2^k a)$, respectively. Suppose $T \in \text{CZO}_{n+\nu}$ satisfies the weak boundedness property. Then:*

(a) *there is a constant $C < \infty$ such that for all $v \in \mathbb{R}^N$*

$$|\langle \phi(\cdot - v), T[\varphi(\cdot - v)] \rangle| \leq C \frac{1+k}{(a2^k)^N} \left(1 + \frac{|w|}{a2^k}\right)^{-N-\nu};$$

(b) *if $T(u^\alpha) = 0$ for all $|\alpha| \leq n$, then there are constants $C < \infty$ and $\delta > 0$ such that for all $v \in \mathbb{R}^N$,*

$$|\langle \phi(\cdot - v), T'[\varphi(\cdot - v)] \rangle| \leq C(a2^k)^{-N-n-\nu} \left(1 + \frac{|w|}{a2^k}\right)^{-N-\delta}.$$

3. THE TRANSLATION LEMMA

This section is devoted to the statement and proof of an estimate for translations in the reflexive spaces $L_p(\ell_q)$. For $q = 2$, this is a square function estimate which was proved, apparently independently, by J. Bourgain [9] (with application to vector-valued singular integrals) and M. Yamazaki [13] (with application to pseudodifferential operators). In fact, Bourgain’s result covers a much more general (and deeper) Banach space -valued situation, but this generality is to a different direction than our present needs. The proof given below is in the same spirit as Yamazaki’s (i.e., vector-valued Calderón–Zygmund theory), but is not an immediate modification since we do not have the Hilbert space structure of ℓ_2 available here.

Lemma 3.1. *Let $1 < p, q < \infty$, and $h_i = 2^i k_i$, where $|k_i| \leq K$ for some $K > 2$ and for all $i \in \mathbb{Z}$. Let $\varphi : \mathbb{R}^N \rightarrow \mathbb{C}$ be a differentiable function with*

$$\|\varphi\|_{L_1} \leq c, \quad |\nabla\varphi(x)| \leq c(1 + |x|)^{-N-1}, \quad c > 0,$$

and denote $\varphi_{2^i} = 2^{-iN}\varphi(2^{-i}\cdot)$ for $i \in \mathbb{Z}$. Then there is a constant $C < \infty$ depending only on p, q, N and c , such that

$$\|(\tau_{h_i}\varphi_{2^i} * f_i)\|_{L_p(\ell_q)} \leq C \log K \|f_i\|_{L_p(\ell_q)}.$$

Here τ_{h_i} denotes the translation operator $f \mapsto f(\cdot - h_i)$.

In particular, if $\text{supp}\widehat{f_i} \subseteq \{|\xi| \leq 2^{-i}\}$ for all $i \in \mathbb{Z}$, then

$$\|(\tau_{h_i}f_i)\|_{L_p(\ell_q)} \leq C \log K \|f_i\|_{L_p(\ell_q)}.$$

Proof. Note first that, once we have proved the first assertion, the second one immediately follows by choosing $\varphi \in \mathcal{S}(\mathbb{R}^N)$ such that $\widehat{\varphi}(\xi) = 1$ for $|\xi| \leq 1$; then $\widehat{\varphi_{2^i}}(\xi) = \widehat{\varphi}(2^i\xi) = 1$ on $\text{supp } \widehat{f_i}$ so that $\varphi_{2^i} * f_i = f_i$. Let us thus consider the first assertion.

For $p = q$, we have $L_q(\ell_q) = \ell_q(L_q)$ and $\|\tau_{h_i}\varphi_{2^i} * f_i\|_{L_q} = \|\varphi_{2^i} * f_i\|_{L_q} \leq \|\varphi_{2^i}\|_{L_1}\|f_i\|_{L_q}$, and $\|\varphi_{2^i}\|_{L_1} = \|\varphi\|_{L_1} \leq c$, so the assertion holds with c in place of $C \log K$. The main part of the proof will consist of verifying that the diagonal-operator-valued kernel $\mathcal{K}(x) = \left(2^{-iN}\varphi(2^{-i}x - k_i)\right)_{i=-\infty}^{\infty}$ of the convolution operator $(f_i) \mapsto (\tau_{h_i}\varphi_{2^i} * f_i)$ satisfies the Hörmander integral condition

$$\int_{|x|>2|y|} \|\mathcal{K}(x - y) - \mathcal{K}(x)\|_{\mathcal{L}(\ell_q)} dx \leq C \log K. \tag{2}$$

This then yields the $L_p(\ell_q)$ boundedness of our operator for all $1 < p < \infty$, with the desired norm estimate, by the well-known vector-valued extension of the theory of singular integrals (see [14]).

For the proof of (2), note first that

$$\begin{aligned} & \|\mathcal{K}(x - y) - \mathcal{K}(x)\|_{\mathcal{L}(\ell_q)} \\ &= \sup_{i \in \mathbb{Z}} 2^{-iN} |\varphi(2^{-i}(x - y) - k_i) - \varphi(2^{-i}x - k_i)| \\ &= \sup_{i \in \mathbb{Z}} 2^{-i(N+1)} \left| \int_0^1 y \cdot \nabla \varphi(2^{-i}(x - \lambda y) - k_i) d\lambda \right| =: \sup_{i \in \mathbb{Z}} \mathcal{K}_i(x, y) \\ &\leq \sup_{2^i \leq |x|/4K} \mathcal{K}_i(x, y) + \sup_{|x|/4K < 2^i < |x|} \mathcal{K}_i(x, y) + \sup_{|x| \leq 2^i} \mathcal{K}_i(x, y). \end{aligned} \tag{3}$$

For $|x| \geq \max(4K \cdot 2^i, 2|y|)$ we have

$$|2^{-i}(x - y) - k_i| \geq 2^{-i}(1 - 2^{-1} - 4^{-1})|x| = 2^{-i-2}|x|,$$

and hence

$$\mathcal{K}_i(x, y) \leq 2^{-i(N+1)}|y|C(2^{-i}|x|)^{-N-1} = C|y| \cdot |x|^{-N-1}.$$

For $|x| \leq 2^i$ we may estimate

$$\mathcal{K}_i(x, y) \leq 2^{-i(N+1)}|y|C \leq C|y| \cdot |x|^{-N-1}.$$

Since

$$\int_{|x|>2|y|} |x|^{-N-1} dx = C|y|^{-1},$$

we have handled the Hörmander estimate of the first and the third term in (3), in fact, with C in place of $C \log K$.

Concerning the second term in (3), we first consider the integral over $|x| > 4K|y|$ and estimate the supremum by the sum

$$\begin{aligned} & \int_{|x|>4K|y|} \sum_{|x|/4K < 2^i < |x|} 2^{-i(N+1)}|y| \int_0^1 |\nabla\varphi(2^{-i}(x - \lambda y) + k_i)|d\lambda dx \\ &= \sum_{2^i > |y|} 2^{-i(N+1)}|y| \int_0^1 \int_{2^i \vee 4K|y| < |x| \leq 4K \cdot 2^i} |\nabla\varphi(2^{-i}(x - \lambda y) + k_i)|dx d\lambda \\ &\leq \sum_{2^i > |y|} 2^{-i}|y| \cdot \|\nabla\varphi\|_{L_1} \leq C, \end{aligned}$$

where in the first equality we could restrict the summation range of the i variable, since for $2^i \leq |y|$ the x integration is over the empty set.

The remaining part in (3) yet to be estimated will give a principal contribution to the estimate, and is in particular responsible for the logarithmic factor $\log K$. As above, we estimate the supremum by the sum and interchange the order of summation and integration, arriving at

$$\begin{aligned} & \sum_{|y|/2K < 2^i < 4K|y|} 2^{-iN} \int |\varphi(2^{-i}(x - y) - k_i) - \varphi(2^{-i}x - k_i)|dx \\ & \leq \sum_{|y|/2K < 2^i < 4K|y|} 2\|\varphi\|_{L_1} \leq C \log K, \end{aligned}$$

where the last estimate is simply counting the number of terms in the sum. This completes the proof of the Hörmander condition (2) and of the Lemma. \square

4. PROOF OF THEOREM 1.1

The proof of Theorem 1.1 is done in three steps.

In the first step we decompose of the operator T . For this purpose, we choose $\Phi \in \mathcal{D}^n(\mathbb{R}^N)$ and $\Psi \in \mathcal{S}(\mathbb{R}^N)$ such that

- Φ is real and radial,
- $\widehat{\Phi}, \widehat{\Psi} \geq 0$,
- $\widehat{\Phi}(u) \geq 1$ for $\frac{1}{2} \leq |u| \leq 2$,
- $\text{supp}\widehat{\Psi} \subseteq \{\frac{1}{2} \leq |u| \leq 2\}$, and
- $\sum_{j \in \mathbb{Z}} \widehat{\Phi}(2^j u) \widehat{\Psi}(2^j u) = 1$ for all $u \in \mathbb{R}^N \setminus \{0\}$.

The existence of such Φ, Ψ is shown in [10] (see also [8]). For $j \in \mathbb{Z}$, we denote $\Phi_{2^j}(u) := 2^{-Nj}\Phi(2^{-j}u)$, $\Psi_{2^j}(u) := 2^{-Nj}\Psi(2^{-j}u)$, $P_j f := \Phi_{2^j} * f$, and $Q_j f = \Psi_{2^j} * f$. Now we can write

$$\langle g, Tf \rangle = \sum_{j,l \in \mathbb{Z}} \langle P_l Q_l g, TP_j Q_j f \rangle = \sum_{j,k \in \mathbb{Z}} \langle Q_{j+k} g, P_{j+k} TP_j Q_j f \rangle,$$

if $f, g \in \mathcal{S}(\mathbb{R}^N)$. The operator $T_{j+k,j} := P_{j+k}TP_j$ is associated with the kernel $K_{j+k,j}$ given by

$$K_{j+k,j}(u, v) = \langle \Phi_{2^{j+k}}(\cdot - u), T[\Phi_{2^j}(\cdot - v)] \rangle.$$

In the second step we fix $k \in \mathbb{Z}$ and estimate the sum over $j \in \mathbb{Z}$ in the above decomposition.

We first consider the case $k \in \mathbb{N}$. For $f, g \in \mathcal{Z}(\mathbb{R}^N)$,

$$\begin{aligned} \left| \sum_{j \in \mathbb{Z}} \langle Q_{j+k}g, T_{j+k,j}Q_jf \rangle \right| &= \left| \sum_{j \in \mathbb{Z}} \langle 2^{(j+k)s}(T_{j+k,j})'Q_{j+k}g, 2^{-(j+k)s}Q_jf \rangle \right| \\ &\leq \left\| \left(\sum_{j \in \mathbb{Z}} |2^{(j+k)s}(T_{j+k,j})'Q_{j+k}g|^{q'} \right)^{1/q'} \right\|_{L_{p'}} \left\| \left(\sum_{j \in \mathbb{Z}} |2^{-(j+k)s}Q_jf|^q \right)^{1/q} \right\|_{L_p}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{p'} = 1$ and $\frac{1}{q} + \frac{1}{q'} = 1$. The second factor is bounded by $C2^{-ks}\|f\|_{\dot{F}_p^{s,q}}$ (cf. [12, 5.1.5]). To estimate the first factor, we note that for $v \in \mathbb{R}^N$

$$\begin{aligned} [(T_{j+k,j})'\phi](v) &= \int_{\mathbb{R}^N} K_{j+k,j}(u, v)\phi(u)du \\ &= \int_{\mathbb{R}^N} 2^{jN}K_{j+k,j}(v + 2^ju, v)\phi(v + 2^ju)du, \end{aligned}$$

and therefore

$$\begin{aligned} A &:= \left\| \left(\sum_{j \in \mathbb{Z}} |2^{(j+k)s}(T_{j+k,j})'Q_{j+k}g|^{q'} \right)^{1/q'} \right\|_{L_{p'}} \\ &\leq \int_{\mathbb{R}^N} \left\| \left(\sum_{j \in \mathbb{Z}} |2^{(j+k)s}2^{jN}K_{j+k,j}(\cdot + 2^ju, \cdot)(Q_{j+k}g)(\cdot + 2^ju)|^{q'} \right)^{1/q'} \right\|_{L_{p'}} du. \end{aligned}$$

Using Lemma 2.1 (a), we obtain the estimate

$$\sup_{v \in \mathbb{R}^N} \sup_{j \in \mathbb{Z}} |2^{jN}K_{j+k,j}(v + 2^ju, v)| \leq C \frac{1+k}{(a2^k)^N} \left(1 + \frac{|u|}{a2^k} \right)^{-N-\nu},$$

where $a > 0$ is such that $\text{supp}\Phi \subseteq \overline{B}(0, a)$. So with Lemma 3.1 we obtain

$$\begin{aligned} A &\leq C \int_{\mathbb{R}^N} \frac{1+k}{(a2^k)^N} \left(1 + \frac{|u|}{a2^k} \right)^{-N-\nu} \log \left(2 + \frac{|u|}{2^k} \right) \|g\|_{\dot{F}_{p'}^{-s,q'}} du \\ &\leq C(1+k)\|g\|_{\dot{F}_{p'}^{-s,q'}}. \end{aligned}$$

Combining this with the bound for the factor f , we have

$$\left| \sum_{j \in \mathbb{Z}} \langle Q_{j+k}g, T_{j+k,j}Q_jf \rangle \right| \leq C(1+k)2^{-ks}\|f\|_{\dot{F}_p^{s,q}}\|g\|_{\dot{F}_{p'}^{-s,q'}}.$$

If k is negative, we proceed in a similar way. Here we use the integral representation for $T_{j+k,j}$ and Lemma 2.1 (b) to obtain

$$\left| \sum_{j \in \mathbb{Z}} \langle Q_{j+k}g, T_{j+k,j}Q_jf \rangle \right| \leq C2^{k(n+\nu-s)} \|f\|_{\dot{F}_p^{s,q}} \|g\|_{\dot{F}_{p'}^{-s,q'}}.$$

As the third step, we carry out the summation over k . By our results from the previous step,

$$\left| \sum_{j,k \in \mathbb{Z}} \langle Q_{j+k}g, T_{j+k,j}Q_jf \rangle \right| \leq C \|f\|_{\dot{F}_p^{s,q}} \|g\|_{\dot{F}_{p'}^{-s,q'}}$$

if $s \in (0, n + \nu)$. Since $\mathcal{Z}(\mathbb{R}^N)$ is dense both in $\dot{F}_p^{s,q}(\mathbb{R}^N)$ and in $\dot{F}_{p'}^{-s,q'}$, the first part of the theorem is proved.

For the second part, we observe that our additional assumptions imply that we can use Lemma 2.1 (b) in Step 2 to show that

$$\left| \sum_{j \in \mathbb{Z}} \langle Q_{j+k}g, T_{j+k,j}Q_jf \rangle \right| \leq C2^{-|k|(n+\nu-|s|)} \|f\|_{\dot{F}_p^{s,q}} \|g\|_{\dot{F}_{p'}^{-s,q'}}.$$

This completes the proof of the theorem.

ACKNOWLEDGEMENTS

Tuomas Hytönen was partially supported by the Finnish Academy of Science and Letters (Vilho, Yrjö and Kalle Väisälä Foundation). Cornelia Kaiser was supported by the Land Baden-Württemberg (Margarete von Wrangell scholarship). The research was carried out while both authors held Visiting Fellow positions at the Centre of Mathematics and its Applications of the Australian National University.

REFERENCES

1. G. DAVID and J.-L. JOURNÉ, A boundedness criterion for generalized Calderón–Zygmund operators, *Ann. of Math. (2)* **120**(1984), No. 2, 371–397.
2. M. FRAZIER, Y.-S. HAN, B. JAWERTH, and G. WEISS, The $T1$ theorem for Triebel–Lizorkin spaces. *Harmonic analysis and partial differential equations (El Escorial, 1987)*, 168–181, *Lecture Notes in Math.*, 1384, Springer, Berlin, 1989.
3. Y.-S. HAN and S. HOFMANN, $T1$ theorems for Besov and Triebel–Lizorkin spaces. *Trans. Amer. Math. Soc.* **337**(1993), No. 2, 839–853.
4. P. G. LEMARIÉ, Continuité sur les espaces de Besov des opérateurs définis par des intégrales singulières. *Ann. Inst. Fourier (Grenoble)* **35**(1985), No. 4, 175–187.
5. R. H. TORRES, Boundedness results for operators with singular kernels on distribution spaces. *Mem. Amer. Math. Soc.* **90**(1991), No. 442, 172 pp.
6. A. YOUSSEFI, Continuité-Besov des opérateurs définis par des intégrales singulières, *Manuscripta Math.* **65**(1989), No. 3, 289–310.
7. T. FIGIEL, Singular integral operators: a martingale approach. *Geometry of Banach spaces (Strobl, 1989)*, 95–110, *London Math. Soc. Lecture Note Ser.*, 158, Cambridge Univ. Press, Cambridge, 1990.

8. T. HYTÖNEN and L. WEIS, A $T1$ theorem for integral transformations with operator-valued kernel, *J. reine angew. Math.* (to appear) [Preprint available at <http://users.utu.fi/~tuohyt/papers.html>]
9. J. BOURGAIN, Vector-valued singular integrals and the H^1 -BMO duality. *Probability theory and harmonic analysis (Cleveland, Ohio, 1983)*, 1–19, *Monogr. Textbooks Pure Appl. Math.*, 98, Dekker, New York, 1986.
10. C. KAISER, Calderón–Zygmund operators with operator-valued kernel on homogeneous Besov spaces. Submitted.
11. C. KAISER, Calderón–Zygmund operators with operator-valued kernel on homogeneous Bessel potential spaces. In preparation.
12. H. TRIEBEL, Theory of function spaces. *Monographs in Mathematics*, vol. 78, Birkhäuser Verlag, Basel, 1983.
13. M. YAMAZAKI, The L^p -boundedness of pseudodifferential operators with estimates of parabolic type and product type. *J. Math. Soc. Japan* **38**(1986), No. 2, 199–225.
14. A. BENEDEK, A.-P. CALDERÓN, and R. PANZONE, Convolution operators on Banach space valued functions, *Proc. Nat. Acad. Sci. U.S.A.* **48**(1962), 356–365.

(Received 29.04.2006)

Authors' addresses:

T. Hytönen
Department of Mathematics
University of Turku
20014 Turku
Finland
E-mail: tuomas.hytonen@utu.fi

C. Kaiser
Mathematisches Institut I
Universität Karlsruhe
Englerstraße 2 76128 Karlsruhe
Germany
E-mail: cornelia.kaiser@math.uni-karlsruhe.de