

Asymptotic Behaviour of a Thin Insulation Problem

Pierpaolo Esposito

*Dipartimento di Matematica, Università di Roma Tor Vergata,
via della Ricerca Scientifica, Roma 00133, Italy*

Giuseppe Riey

*Dipartimento di Matematica, Università di Roma Tor Vergata,
via della Ricerca Scientifica, Roma 00133, Italy*

Received June 10, 2002

Revised manuscript received September 10, 2002

We consider a problem of thermal insulation and we study, by means of Γ -convergence, the best way to put the insulator when its mass tends to zero.

Keywords: Γ -convergence, thermal insulation, asymptotic behaviour

2000 Mathematics Subject Classification: 35B40, 35J05, 35B37

1. Introduction

Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a $C^{1,1}$ bounded open set. For any point $\sigma \in \partial\Omega$, we denote by $n(\sigma)$ the unit outer normal to $\partial\Omega$. We cover externally the boundary of the region Ω by a thin layer Σ_ε of material of small conductivity $\delta > 0$ as follows. For any $h \in C^0(\partial\Omega, [0, +\infty[)$, we define

$$\Sigma_\varepsilon = \Sigma_\varepsilon(h) := \{x \in \mathbb{R}^n : x = \sigma + th(\sigma)n(\sigma), \sigma \in \partial\Omega, t \in (0, \varepsilon)\},$$

where $\varepsilon \in \mathbb{R}^+$ is small enough in a way to obtain $\overline{\Sigma_\varepsilon} \cap \overline{\Omega} = \partial\Omega$. Set

$$\Omega_\varepsilon := \overline{\Omega} \cup \Sigma_\varepsilon$$

and let $a_{\varepsilon,\delta} : \Omega_\varepsilon \rightarrow \mathbb{R}$ be the function representing the conductivity of Ω_ε , defined as

$$a_{\varepsilon,\delta}(x) := \begin{cases} 1 & \text{if } x \in \overline{\Omega} \\ \delta & \text{if } x \in \Sigma_\varepsilon. \end{cases}$$

Denoting by $u(x)$ the temperature at the point x (which is supposed to be identically equal to 0 out of Ω_ε), then $u(x)$ satisfies the nonhomogeneous equation

$$\begin{cases} u \in H_0^1(\Omega_\varepsilon) \\ -\operatorname{div}(a_{\varepsilon,\delta}\nabla u) = f \quad \text{in } \mathcal{D}'(\Omega_\varepsilon) \end{cases} \quad (1)$$

where $f \in L^p(\Omega)$, $p > \frac{2n}{n+2}$, represents a given heat source (f is extended to 0 out of Ω). Equivalently, u is a critical point of the functional $F_{\varepsilon,\delta} : L^1(\mathbb{R}^n) \rightarrow \mathbb{R} \cup \{+\infty\}$ defined by:

$$F_{\varepsilon,\delta}(u) := \begin{cases} \frac{1}{2} \int_\Omega |\nabla u|^2 + \frac{\delta}{2} \int_{\Sigma_\varepsilon} |\nabla u|^2 - \int_\Omega f u & \text{if } u \in H_0^1(\Omega_\varepsilon) \\ +\infty & \text{elsewhere.} \end{cases}$$

Let us recall what it is known on the asymptotic behaviour of problem (1) as $(\varepsilon, \delta) \rightarrow (0, 0)$. It is intuitively clear that, if $\varepsilon \ll \delta$, we may neglect the insulator and thus we expect to obtain in the limit the usual Dirichlet problem. If instead $\delta \ll \varepsilon$, we expect to have not heat transmission through $\partial\Omega$ and thus to obtain the Neumann problem.

The most interesting case is when $\varepsilon \approx \delta$, that is $\varepsilon = k\delta$. Up to a renormalization we can assume $k = 1$ and set $F_\varepsilon := F_{\varepsilon,\varepsilon}$. Results due to Acerbi and Buttazzo ([1],[7]) imply the following theorem (see also [3], [4], [6] and [8]), which is the starting point of our work.

Theorem 1.1. *For any $h \in C^0(\partial\Omega; [0, +\infty[)$, the sequence $\{F_\varepsilon\}$ Γ -converges with respect to the $L^1(\mathbb{R}^n)$ -topology to the functional $F(h, \cdot) : L^1(\mathbb{R}^n) \rightarrow [0, +\infty]$ defined by:*

$$F(h, u) := \begin{cases} \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{2} \int_{\partial\Omega} \frac{u^2}{h} - \int_{\Omega} f u & \text{if } u \in H^1(\Omega) \\ +\infty & \text{elsewhere.} \end{cases} \tag{2}$$

Let $m > 0$; define Ξ_m as the set of all measurable functions $h : \partial\Omega \rightarrow [0, +\infty[$ such that $\int_{\partial\Omega} h = m$. In the sequel of the paper, the functional $F(h, u)$ will have the same expression also for functions $h \in \Xi_m$ which are not continuous in general.

Let (\bar{h}_m, u_m) denote the pair which minimizes F in $\Xi_m \times H^1(\Omega)$ and set $h_m := \frac{\bar{h}_m}{m}$. The purpose of the present paper is to study, by means of Γ -convergence, the asymptotic behaviour of h_m as m goes to 0.

Let us suppose from now on $p > n$. For fixed $h \in \Xi_m$, $F(h, u)$ is strictly convex in u and therefore it admits a unique minimizer $u^h \in H^1(\Omega)$, which satisfies the following equation:

$$\begin{cases} -\Delta u^h(x) = f(x) & x \in \Omega \\ h \frac{\partial u^h}{\partial n}(x) + u^h(x) = 0 & x \in \partial\Omega. \end{cases} \tag{3}$$

The aim now is to maximize the weighted average of the temperature $\int_{\Omega} f u$. If $f \geq 0$, this means to disperse heat as less as possible. This can be seen as a problem of optimal control (where h represents the control variable). A direct computation shows that

$$F(h, u^h) = -\frac{1}{2} \int_{\Omega} f u^h$$

and so we are led to consider the following problem:

$$\min_{h \in \Xi_m} \min_{u \in H^1(\Omega)} F(h, u). \tag{4}$$

It can be seen (see Section 3) that u_m is the unique solution of the problem

$$\min_{u \in H^1(\Omega)} \left\{ \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{2m} \left(\int_{\partial\Omega} |u| \right)^2 - \int_{\Omega} f u \right\}$$

and

$$h_m = \frac{|\int_{\partial\Omega} u_m|}{\int_{\partial\Omega} |u_m|}.$$

Before stating our main result, we first recall that, from classical elliptic theory and Sobolev embeddings (see for instance [5]), there exists an unique solution $\bar{u} \in C^{1,1-\frac{2}{p}}(\bar{\Omega}) \cap W^{2,p}(\Omega)$ of the following equation:

$$\begin{cases} -\Delta u(x) = f(x) & x \in \Omega \\ u(x) = 0 & x \in \partial\Omega. \end{cases} \tag{5}$$

We remark that the function \bar{u} can be regarded as the temperature in case of no insulation. Let us denote

$$M = \max_{\partial\Omega} \left| \frac{\partial\bar{u}}{\partial n} \right|, \quad K^\pm := \{x \in \partial\Omega : \frac{\partial\bar{u}}{\partial n}(x) = \pm M\} \tag{6}$$

and define

$$v_m := \frac{u_m - \bar{u}}{m}.$$

Our main result is stated in the following

Theorem 1.2. *Up to a subsequence, $\left\{ \frac{v_m}{\int_{\partial\Omega} |v_m|} \right\}$ weakly* converges in $\mathcal{M}(\partial\Omega)$ (see Definition 2.1) as m goes to 0 to a measure λ which satisfies*

$$\text{spt } \lambda \subseteq K^+ \cup K^- \tag{7}$$

$$\lambda \text{ is positive (resp. negative) in } K^- \text{ (resp. } K^+) \tag{8}$$

$$\int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda = -M. \tag{9}$$

Moreover

$$\lim_{m \rightarrow 0} \int_{\partial\Omega} |v_m| = |\lambda|(\partial\Omega) = M,$$

where $|\lambda|$ denotes the total variation of λ .

Theorem 1.2 and the fact that $h_m = \frac{|v_m|}{\int_{\partial\Omega} |v_m|}$ (see (11) below) imply the following result.

Theorem 1.3. *The sequence $\{h_m\}$ weakly* converges in $\mathcal{M}(\partial\Omega)$ as m goes to 0 to the total variation of λ .*

Theorem 1.3 states that $\{h_m\}$ converges to a measure concentrated on $K^+ \cup K^-$; physically this means that, when m is infinitesimal, the insulator has to be put in the points where the dispersion is maximal.

2. Notation and preliminaries

We first introduce some notation and briefly recall some basic results about Γ -convergence and Measure Theory.

Let Y be a metric space locally compact and separable. We denote by $\mathcal{M}(Y)$ the space of all finite and real Radon measures on Y and for any $\mu \in \mathcal{M}(Y)$ we denote by $|\mu|$ the total variation of μ .

Definition 2.1 (Weakly* convergence). Let $\mu \in \mathcal{M}(Y)$ and let $\{\mu_n\} \subset \mathcal{M}(Y)$. We say that $\{\mu_n\}$ weakly* converges to μ if

$$\lim_{n \rightarrow +\infty} \int_Y u \, d\mu_n = \int_Y u \, d\mu \quad \text{for every } u \in C_0(Y).$$

Theorem 2.2. Let $\{\mu_n\} \subset \mathcal{M}(Y)$ be a sequence such that $\sup_n |\mu_n|(Y) < +\infty$. Then $\{\mu_n\}$ has a weakly* converging subsequence. Moreover the map $\mu \mapsto |\mu|(Y)$ is lower semicontinuous with respect to the weak* convergence.

Remark 2.3. The topology induced by the weak* convergence defined above is metrizable on the bounded sets of $\mathcal{M}(Y)$.

For more details about this topic see for instance [2].

Definition 2.4. Let X be a topological space and let $F_n, F : X \rightarrow \mathbb{R} \cup \{+\infty\}$. Define the Γ -lim inf and the Γ -lim sup of F_n as:

$$\begin{aligned} \Gamma\text{-lim inf } F_n(x) &:= \inf \left\{ \liminf_{n \rightarrow \infty} F_n(x_n) : x_n \rightarrow x \right\} \\ \Gamma\text{-lim sup } F_n(x) &:= \inf \left\{ \limsup_{n \rightarrow \infty} F_n(x_n) : x_n \rightarrow x \right\}. \end{aligned}$$

We say that F_n Γ -converges to F if for all $x \in X$ we have:

$$\Gamma\text{-lim inf}_{n \rightarrow \infty} F_n(x) = \Gamma\text{-lim sup}_{n \rightarrow \infty} F_n(x) = F(x).$$

Definition 2.5. A sequence of functionals $F_n : X \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be sequentially equicoercive if, for any sequence $\{x_n\}$ such that $\sup_n F_n(x_n) < +\infty$, there exists a convergent subsequence.

Theorem 2.6. Let $\{F_n\}$ be a sequence of sequentially equicoercive functionals defined on X and Γ -converging to F . Then there exists $\min_X F$ and $\min_X F = \liminf_{n \rightarrow \infty} \min_X F_n$. Moreover, if x_n is a minimizer of F_n , then every limit of a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ is a minimizer of F .

For more details about this topic see for instance [9].

In the sequel we identify any $f \in L^1(\Omega)$ (resp. $f \in L^1(\partial\Omega)$) with the measure $f \, dx$ (resp. $f \, d\sigma$).

3. Main results

In order to study $\min_{h \in \Xi_m} \min_{u \in H^1(\Omega)} F(h, u)$, according to [7], we consider the equivalent problem:

$$\min_{u \in H^1(\Omega)} \min_{h \in \Xi_m} F(h, u). \tag{10}$$

Denoting by $\bar{h}_m(u) \in \Xi_m$ the unique solution of the problem

$$\min_{h \in \Xi_m} F(h, u),$$

it is possible to see that

$$\bar{h}_m(u) = m \frac{|u|}{\int_{\partial\Omega} |u|}. \tag{11}$$

Thus, setting

$$E_m(u) := F(\bar{h}_m(u), u) = \frac{1}{2} \int_{\Omega} |\nabla u|^2 + \frac{1}{2m} \left(\int_{\partial\Omega} |u| \right)^2 - \int_{\Omega} f u,$$

problem (10) becomes:

$$\min_{u \in H^1(\Omega)} E_m(u). \tag{12}$$

It is possible to prove (see [7]) that $E_m(u)$ is strictly convex, coercive and lower semicontinuous with respect to the weak topology of $H^1(\Omega)$.

Hence it immediately follows:

Theorem 3.1. *Problem (12) admits a unique solution $u_m \in H^1(\Omega)$, which satisfies*

$$\begin{cases} -\Delta u(x) = f(x) & x \in \Omega \\ \frac{\partial u}{\partial n}(x) + \frac{1}{m} \text{sign}(u(x)) \int_{\partial\Omega} |u| \ni 0 & x \in \partial\Omega. \end{cases} \tag{13}$$

In (13) the map sign is defined as $\text{sign}(t) := \pm 1$ if $t \gtrless 0$ and $\text{sign}(0) = [-1, 1]$.

Corollary 3.2. *Let $\bar{h}_m(u)$ be defined as in (11) and let u_m be given by Theorem 3.1. Then the pair $(\bar{h}_m(u_m), u_m)$ solves the minimization problem (10) or equivalently (4).*

Defining $h_m := \frac{\bar{h}_m(u_m)}{m}$ and recalling (11), we have $\int_{\partial\Omega} h_m = 1$ and $h_m \geq 0$. Hence by Theorem 2.2 the sequence $\{h_m\}$ admits a subsequence weakly* converging in $\mathcal{M}(\partial\Omega)$ as $m \rightarrow 0$.

In the case of Ω an annular region in \mathbb{R}^2 and $f \equiv 1$, solving explicitly equation (13) in polar coordinates, it turns out that the optimal way to insulate Ω is achieved by putting the insulator on the inner circle, the set of boundary points with negative mean curvature, see [7] for details. This could suggest that h_m tends, as $m \rightarrow 0$, to a measure concentrated where the mean curvature of $\partial\Omega$ relative to Ω is minimal.

Nevertheless, this is false, as it follows from the following example.

Example 3.3. Let $f \equiv 1$ and $\Omega = (B_R(0) \setminus \overline{B_r(0)}) \cup B_\rho(p) \subset \mathbb{R}^2$ for $R > r > 0$, $\rho > 0$ and $|p| > R + \rho$. We define u_m as

$$u_m(x) = \begin{cases} -\frac{1}{4} (|x|^2 - r^2) + \frac{(R^2 - r^2)}{4 \log \frac{R}{r}} \log \frac{|x|}{r} & \text{if } x \in B_R(0) \setminus \overline{B_r(0)} \\ \frac{m}{4\pi} + \frac{\rho^2 - |x-p|^2}{4} & \text{if } x \in B_\rho(p). \end{cases}$$

It can be seen that $u_m |_{\partial B_r(0)} = u_m |_{\partial B_R(0)} \equiv 0$, $u_m |_{\partial B_\rho(p)} \equiv \frac{m}{4\pi}$ and

$$\frac{\partial u_m}{\partial n}(x) = \begin{cases} \frac{1}{2}r - \frac{R^2-r^2}{4r \log \frac{R}{r}} & \text{if } x \in \partial B_r(0) \\ -\frac{1}{2}R + \frac{R^2-r^2}{4R \log \frac{R}{r}} & \text{if } x \in \partial B_R(0) \\ -\frac{1}{2}\rho & \text{if } x \in \partial B_\rho(p). \end{cases}$$

Hence $\int_{\partial\Omega} |u_m| = \frac{m}{2}\rho$ and

$$0 \in \frac{\partial u_m}{\partial n}(x) + \frac{1}{m} \text{sign}(u(x)) \int_{\partial\Omega} |u_m|$$

if and only if $\rho \geq \rho_0 = \rho_0(R, r) = \max \left\{ \left| R - \frac{R^2-r^2}{2R \log \frac{R}{r}} \right|, \left| r - \frac{R^2-r^2}{2r \log \frac{R}{r}} \right| \right\}$.

For fixed r, R , $\rho \geq \rho_0$, we can choose a point p so that $|p| > R + \rho$: so u_m turns out to be the unique solution of equation (13) and $h_m = \frac{|u_m|}{\int_{\partial\Omega} |u_m|} \in \Xi_m$ satisfies $\text{spt } h_m = \partial B_\rho(p)$ for any $m > 0$.

The set of boundary points with minimal mean curvature is $B_r(0)$ and it is different from $\text{spt } h_m$. According to Theorems 1.2 and 1.3, it is easy to check that $\partial B_\rho(p) \subseteq K^+ \cup K^-$ and the equality holds if $\rho > \rho_0$.

We observe that, if we write $u \in H^1(\Omega)$ in the form

$$u = \bar{u} + m v, \quad \bar{u} \text{ the solution of (5), } v \in H^1(\Omega)$$

and if $I, \tilde{J}_m : H^1(\Omega) \rightarrow [0, +\infty[$ are the functionals defined as

$$\begin{cases} I(u) := \frac{1}{2} \int_{\Omega} |\nabla u|^2 - \int_{\Omega} f u \\ \tilde{J}_m(v) := \frac{m}{2} \int_{\Omega} |\nabla v|^2 + \frac{1}{2} \left(\int_{\partial\Omega} |v| \right)^2 + \int_{\partial\Omega} \frac{\partial \bar{u}}{\partial n} v, \end{cases} \tag{14}$$

then we have:

$$E_m(\bar{u} + m v) = I(\bar{u}) + m \tilde{J}_m(v). \tag{15}$$

In particular we can write the function u_m of Theorem 3.1 in the form

$$u_m = \bar{u} + m v_m,$$

where v_m is the unique minimizer of \tilde{J}_m and satisfies:

$$\begin{cases} \Delta v_m(x) = 0 & x \in \Omega \\ \text{sign}(v_m(x)) \int_{\partial\Omega} |v_m(x)| + m \frac{\partial v_m(x)}{\partial n} + \frac{\partial \bar{u}(x)}{\partial n} \ni 0 & x \in \partial\Omega. \end{cases} \tag{16}$$

Since \bar{u} vanishes on $\partial\Omega$ we have

$$h_m = \frac{|u_m|}{\int_{\partial\Omega} |u_m|} = \frac{|v_m|}{\int_{\partial\Omega} |v_m|}.$$

We now focus the attention on the asymptotic behaviour of the functions v_m .

For any $\varphi \in H^{\frac{1}{2}}(\partial\Omega)$, there exists a unique function $v_\varphi \in H^1(\Omega)$ such that:

$$\int_{\Omega} |\nabla v_\varphi|^2 = \inf_{\{w \in H^1(\Omega), w|_{\partial\Omega} = \varphi\}} \int_{\Omega} |\nabla w|^2. \tag{17}$$

So we can define a new functional $J_m : \mathcal{M}(\partial\Omega) \rightarrow \mathbb{R}$ as:

$$J_m(\varphi) := \begin{cases} \frac{m}{2} \int_{\Omega} |\nabla v_\varphi|^2 + \frac{1}{2} \left(\int_{\partial\Omega} |\varphi| \right)^2 + \int_{\partial\Omega} \frac{\partial \bar{u}}{\partial n} \varphi & \text{if } \varphi \in H^{\frac{1}{2}}(\partial\Omega) \\ +\infty & \text{elsewhere.} \end{cases}$$

Notice that $v_m|_{\partial\Omega}$ is the unique minimizer of J_m . We now give a result of compactness and Γ -convergence for the functionals J_m .

Theorem 3.4 (Compactness for J_m). *Let $\{\varphi_m\} \subset H^{\frac{1}{2}}(\partial\Omega)$ be such that $\sup_m J_m(\varphi_m) < +\infty$. Then there exist $\mu \in \mathcal{M}(\partial\Omega)$ and a subsequence of $\{\varphi_m\}$ weakly* converging to μ in $\mathcal{M}(\partial\Omega)$ as $m \rightarrow 0$.*

Proof. Setting $M := \max_{x \in \partial\Omega} \left| \frac{\partial \bar{u}}{\partial n}(x) \right|$ and $T := \sup_m J_m(\varphi_m) < +\infty$, we have:

$$M \int_{\partial\Omega} |\varphi_m| - 2M^2 \leq \frac{1}{2} \left(\int_{\partial\Omega} |\varphi_m| \right)^2 - M \int_{\partial\Omega} |\varphi_m| \leq T$$

and hence $\int_{\partial\Omega} |\varphi_m| \leq \frac{T}{M} + 2M$. The thesis now follows from Theorem 2.2. □

Theorem 3.5 (Γ -convergence for J_m). *The sequence $\{J_m\}$ Γ -converges, with respect to the weak* convergence on $\mathcal{M}(\partial\Omega)$, as $m \rightarrow 0$, to the functional $J : \mathcal{M}(\partial\Omega) \rightarrow \mathbb{R}$ defined as*

$$J(\mu) := \frac{1}{2} (|\mu|(\partial\Omega))^2 + \int_{\partial\Omega} \frac{\partial \bar{u}}{\partial n} d\mu.$$

Proof. The Γ -lim inf inequality follows from the lower semicontinuity of the total variation with respect to the weak* convergence in $\mathcal{M}(\partial\Omega)$.

To prove the Γ -lim sup inequality, it is enough to find, for any $\mu \in \mathcal{M}(\partial\Omega)$, a sequence $w_m \in H^1(\Omega)$ such that $\varphi_m := w_m|_{\partial\Omega}$ weakly* converges to μ in $\mathcal{M}(\partial\Omega)$ and $\limsup_{m \rightarrow 0} \tilde{J}_m(w_m) \leq J(\mu)$, because, in view of (17),

$$J_m(\varphi_m) = \tilde{J}_m(v_{\varphi_m}) \leq \tilde{J}_m(w_m) \quad \forall m.$$

To this aim we make the following construction. Set $D := \{x \in \mathbb{R}^{n-1} : |x| < 1\}$, let $\{S_i, r_i : D \rightarrow S_i\}_{i=1, \dots, N}$ be a local coordinates system and let $\{\psi_i\}_{i=1, \dots, N}$ be a unit partition subordinate to the family $\{S_i\}$. Let $\mu_i \in \mathcal{M}(D)$ be defined as the push-forward of $\psi_i \mu$ through the map r_i^{-1} , $\mu_i := (r_i^{-1})_{\#} [\psi_i \mu]$.

Since $\text{spt } \mu_i \subset D$, by Theorem 2.2 in [2] there exists a sequence $\{\tilde{\mu}_i^\varepsilon\} \subset C_0^\infty(D)$ such that:

$$\tilde{\mu}_i^\varepsilon \text{ weakly* converges to } \mu_i \tag{18}$$

$$|\tilde{\mu}_i^\varepsilon|(D) \rightarrow |\mu_i|(D) \tag{19}$$

$$|\tilde{\mu}_i^\varepsilon| = O\left(\frac{1}{\varepsilon^{n-1}}\right), \quad |\nabla \tilde{\mu}_i^\varepsilon| = O\left(\frac{1}{\varepsilon^n}\right). \tag{20}$$

Let $\mu_i^\varepsilon \in \mathcal{M}(\partial\Omega)$ be defined as the push-forward of $\tilde{\mu}_i^\varepsilon$ through the map r_i and let us identify the measure μ_i^ε with the function $\mu_i^\varepsilon := \tilde{\mu}_i^\varepsilon \circ r_i^{-1} \in C_0^\infty(S_i)$. Set $\mu_\varepsilon := \sum_{i=1}^N \mu_i^\varepsilon$.

By construction we get that, as $\varepsilon \rightarrow 0$,

$$\mu_\varepsilon \text{ weakly* converges to } \mu \text{ in } \mathcal{M}(\partial\Omega) \tag{21}$$

and

$$|\mu_\varepsilon|(\partial\Omega) \rightarrow |\mu|(\partial\Omega). \tag{22}$$

We now want to extend μ_ε in the interior of Ω . Set $d(x) := \text{dist}(x, \partial\Omega)$ and for $\alpha > 0$ define $U := \{x \in \Omega : d(x) < \alpha\}$ and $\tilde{U} := \{x \in U : d(x) \in (0, \frac{\alpha}{2})\}$. For α small enough, the orthogonal projection on $\partial\Omega$, $\pi : U \rightarrow \partial\Omega$ is well defined and let $\gamma_\varepsilon : [0, \alpha] \times \partial\Omega \rightarrow \mathbb{R}$ be the function defined as:

$$\gamma_\varepsilon(t, y) := \begin{cases} \mu_\varepsilon(y) & \text{if } t \in [0, \frac{\alpha}{2}] \\ \frac{2(\alpha-t)}{\alpha} \mu_\varepsilon(y) & \text{if } t \in (\frac{\alpha}{2}, \alpha]. \end{cases}$$

Setting

$$w_m(x) := \begin{cases} \gamma_{m^{1/(4n)}}(d(x), \pi(x)) & x \in U \\ 0 & x \in \Omega \setminus U, \end{cases}$$

it is easy to check that $w_m \in H^1(\Omega)$. Moreover, using (20), the lipschitz property of the map π and the fact that $|\nabla d(x)| = 1$ a.e., we infer the following estimate:

$$\begin{aligned} |\nabla w_{m^{1/(4n)}}(x)| &\leq \left| \frac{\partial \gamma_{m^{1/(4n)}}}{\partial t}(d(x), \pi(x)) \right| |\nabla d(x)| \\ &\quad + \left| \frac{\partial \gamma_{m^{1/(4n)}}}{\partial y}(d(x), \pi(x)) \right| |\nabla \pi(x)| \\ &\leq C \left[\left| \frac{\partial \gamma_{m^{1/(4n)}}}{\partial t}(d(x), \pi(x)) \right| + \left| \frac{\partial \gamma_{m^{1/(4n)}}}{\partial y}(d(x), \pi(x)) \right| \right] \\ &= \begin{cases} C |\nabla_\tau \mu_{m^{1/(4n)}}(\pi(x))| & x \in \tilde{U} \\ C \left| -\frac{2}{\alpha} \mu_{m^{1/(4n)}}(\pi(x)) + \frac{2(\alpha-d(x))}{\alpha} \nabla_\tau \mu_{m^{1/(4n)}}(\pi(x)) \right| & x \in U \setminus \tilde{U} \end{cases} \\ &= O\left(\frac{1}{m^{1/4}}\right) \end{aligned}$$

for suitable $C > 0$. Therefore we get

$$m \int_\Omega |\nabla w_{m^{1/(4n)}}|^2 \leq \tilde{C} \sqrt{m} \rightarrow 0 \text{ as } m \rightarrow 0$$

for some $\tilde{C} > 0$.

This concludes the proof of the Γ -limsup inequality. □

In order to characterize the properties of minimizers of J , we write it in a slightly different way. Set $\mathcal{M}_1(\partial\Omega) := \{\mu \in \mathcal{M}(\partial\Omega) : |\mu|(\partial\Omega) = 1\}$ and consider $\tilde{J} : [0, +\infty) \times \mathcal{M}_1(\partial\Omega) \rightarrow \mathbb{R}$ defined as:

$$\tilde{J}(t, \lambda) := J(t\lambda) = \frac{1}{2}t^2 + t \int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda. \tag{23}$$

We have:

$$\min_{\mu \in \mathcal{M}(\partial\Omega)} J(\mu) = \min_{\lambda \in \mathcal{M}_1(\partial\Omega)} \min_{t \geq 0} \tilde{J}(t, \lambda). \tag{24}$$

An easy computation shows that

$$\min_{t \geq 0} \tilde{J}(t, \lambda) = \begin{cases} 0 & \text{if } \int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda \geq 0 \\ -\frac{1}{2} \left(\int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda \right)^2 & \text{if } \int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda < 0 \end{cases}$$

and hence, denoted by $(\bar{t}, \bar{\lambda})$ a pair which minimizes \tilde{J} , we have that:

$$\bar{t} = - \int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\bar{\lambda} > 0 \text{ and } \bar{\lambda} \text{ minimizes } \lambda \mapsto \int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda. \tag{25}$$

We have that $\int_{\partial\Omega} \frac{\partial\bar{u}}{\partial n} d\lambda \geq -M$ and the equality holds if and only if

$$\text{spt } \bar{\lambda} \subseteq K^+ \cup K^-, \quad \bar{\lambda} \geq 0 \text{ (resp. } \leq 0) \text{ in } K^- \text{ (resp. } K^+), \quad \bar{t} = M, \tag{26}$$

where M and K^\pm are defined in (6). Finally by Theorems 2.6, 3.4, 3.5 we can deduce the proof of Theorem 1.2.

Proof of Theorem 1.2. Since $\sup_m J_m(v_m|_{\partial\Omega}) \leq \sup_m J_m(0) = 0 < +\infty$, up to a subsequence, $v_m|_{\partial\Omega}$ weakly* converges in $\mathcal{M}(\partial\Omega)$ to a minimum point μ of J which must have the form $\mu = t\lambda$ with (t, λ) satisfying (25).

Hence, since the total variation is lower semicontinuous with respect to the weak* convergence,

$$|\mu|(\partial\Omega) = M \leq \liminf_{m \rightarrow 0} \int_{\partial\Omega} |v_m|.$$

For any converging subsequence of $\int_{\partial\Omega} |v_m|$ with limit s we have $s \geq M$ and, along this subsequence,

$$-\frac{1}{2}M^2 = J(\mu) = \lim_{m \rightarrow 0} J_m(v_m|_{\partial\Omega}) \geq \frac{1}{2}s^2 - Ms.$$

Since $\frac{1}{2}t^2 - Mt > -\frac{1}{2}M^2$ for $t > M$, then $s = M$ for any converging subsequence of $\int_{\partial\Omega} |v_m|$ and this implies

$$\lim_{m \rightarrow 0} \int_{\partial\Omega} |v_m| = M.$$

Hence $\frac{v_m}{\int_{\partial\Omega} |v_m|}$ weakly* converges to the measure λ in $\mathcal{M}(\partial\Omega)$ and λ satisfies (26). \square

Proof of Theorem 1.3. Up to a subsequence, $h_m \in \mathcal{M}_1(\partial\Omega)$ weakly* converges to some ν in $\mathcal{M}(\partial\Omega)$ with the properties $|\lambda| \leq \nu$ and $\nu(\partial\Omega) \leq 1 = |\lambda|(\partial\Omega)$. Hence $\nu = |\lambda|$ and the proof is completed. \square

Acknowledgements. This paper is the outcome of some discussions held in Cortona on Summer 2001, during a School on Calculus of Variations. We would thank the other students of the school for the nice company, Professors Giuseppe Buttazzo and Bernard Dacorogna for their very interesting lectures and the SMI for the organization of this event.

In particular, we wish to express our deep gratitude to Professor Buttazzo for his helpful advice and comments about this paper.

References

- [1] E. Acerbi, G. Buttazzo: Reinforcement problems in the calculus of variations, *Ann. Inst. H. Poincaré - Analyse Non Linéaire* 3(4) (1986) 273–284.
- [2] L. Ambrosio, N. Fusco, D. Pallara: *Functions of Bounded Variation and Free Discontinuity Problems*, Clarendon Press, Oxford (2000).
- [3] M. Boutkrida, J. Mossino, G. Moussa: On the torsional rigidity problem with nonhomogeneous reinforcement, *Papers in memory of Ennio De Giorgi, Ricerche Mat.* 48, suppl. (1999) 1–24.
- [4] M. Boutkrida, J. Mossino, G. Moussa: On nonhomogeneous reinforcements of varying shape and different exponents, *Boll. U.M.I. Sez. B Artic. Ric. Mat.* (8) 2(3) (1999) 517–536.
- [5] H. Brezis: *Analyse Fonctionnelle - Théorie et Applications*, Masson, Paris (1983).
- [6] H. Brezis, L. A. Caffarelli, A. Friedman: Reinforcement problems for elliptic equations and variational inequalities, *Ann. Mat. Pura Appl.* 123(4) (1980) 219–246.
- [7] G. Buttazzo: An optimization problem for thin insulating layers around a conducting medium, in: *Boundary Control and Boundary Variations (Nice, 1986)*, *Lecture Notes in Comput. Sci.* 100, Springer (1988) 91–95.
- [8] G. Buttazzo: Thin insulating layers: The optimization point of view, in: *Proceedings of "Material Instabilities in Continuum Mechanics and Related Mathematical Problems"*, Edinburgh 1985-1986, Oxford University Press, Oxford (1988) 11-19.
- [9] G. Dal Maso: *An Introduction to Γ -Convergence*, Birkhäuser, Boston (1993).