

M-Structures in Vector-Valued Polynomial Spaces*

Verónica Dimant

*Departamento de Matemática, Universidad de San Andrés,
Vito Dumas 284, (B1644BID) Victoria, Buenos Aires, Argentina and CONICET
vero@udesa.edu.ar*

Silvia Lassalle†

*Departamento de Matemática - Pab I, Facultad de Cs. Exactas y Naturales,
Universidad de Buenos Aires, (1428) Buenos Aires, Argentina and CONICET
slassall@dm.uba.ar*

Received: February 18, 2011

Revised manuscript received: November 07, 2011

This paper is concerned with the study of M -structures in spaces of polynomials. More precisely, we discuss for E and F Banach spaces, whether the class of weakly continuous on bounded sets n -homogeneous polynomials, $\mathcal{P}_w(^nE, F)$, is an M -ideal in the space of continuous n -homogeneous polynomials $\mathcal{P}(^nE, F)$. We show that there is some hope for this to happen only for a finite range of values of n . We establish sufficient conditions under which the problem has positive and negative answers and use the obtained results to study the particular cases when $E = \ell_p$ and $F = \ell_q$ or F is a Lorentz sequence space $d(w, q)$. We extend to our setting the notion of property (M) introduced by Kalton which allows us to lift M -structures from the linear to the vector-valued polynomial context. Also, when $\mathcal{P}_w(^nE, F)$ is an M -ideal in $\mathcal{P}(^nE, F)$ we prove a Bishop-Phelps type result for vector-valued polynomials and relate norm-attaining polynomials with farthest points and remotal sets.

Keywords: M -ideals, homogeneous polynomials, weakly continuous on bounded sets polynomials

2000 Mathematics Subject Classification: 47H60, 46B04, 47L22, 46B20

Introduction

M -ideals emerged in the geometric theory of Banach spaces as a generalization, to the Banach space setting, of the closed two-sided ideals in a C^* -algebra. This notion, introduced by Alfsen and Effros in their seminal article [3] of 1972, leads us to a better understanding of the isometric structure of a Banach space in terms of geometric and analytic properties of the closed unit ball of the dual space. To be more precise, a closed subspace J of a Banach space X is an M -ideal in X , if its annihilator, J^\perp , is the kernel of a projection P on the dual space X^* such that $\|x^*\| = \|P(x^*)\| + \|x^* - P(x^*)\|$, for all $x^* \in X^*$. When J is an M -ideal in X , the canonical complement of J^\perp in X^* is (isometrically) identified with J^* . Then, we

*Partially supported by CONICET-PIP 11220090100624.

†The second author was also partially supported by UBACyT X218 and UBACyT X038.

may write $X^* = J^\perp \oplus_1 J^*$, which in some sense tells us that there is a maximum norm structure underlying the geometry of the unit ball of X and this structure is closely related to J . If it is possible to decompose X as $J \oplus_\infty \tilde{J}$, for some closed subspace \tilde{J} of X , we say that J is an M -**summand** of X . Clearly, M -summands are M -ideals, but there exist subtle differences. For instance, c_0 is an M -ideal in ℓ_∞ and it is not an M -summand. Since M -ideals appeared, they have been intensively studied. A comprehensive exposition of the main developments in this subject can be found in the outstanding book by Hardmand, Werner and Werner [24].

The Gelfand-Naimark theorem states that any arbitrary C^* -algebra is isometrically $*$ -isomorphic to a C^* -algebra of bounded operators on a Hilbert space. Here the only norm closed two-sided $*$ -ideal is the subspace of compact operators. Then, it is natural to investigate under which conditions the closed subspace J of compact operators between Banach spaces E and F , $J = \mathcal{K}(E, F)$, results an M -ideal in $X = \mathcal{L}(E, F)$, the space of linear and bounded operators, endowed with the supremum norm. During the last thirty years a number of papers have been devoted to this question (see, for example [24, 25, 26, 27, 28, 30, 31]), where the case $E = F$ is of special interest.

In this paper we focus our study in determining the presence of an M -structure in the space of continuous n -homogeneous polynomials between Banach spaces E and F , denoted by $\mathcal{P}(^n E, F)$. Here the lack of linearity and, more specifically, the degree of homogeneity will play a crucial role. In the polynomial setting, the space of *compact operators* is usually replaced by the space of *homogeneous polynomials which are weakly continuous on bounded sets*, denoted by $\mathcal{P}_w(^n E, F)$. Recall that a polynomial $P \in \mathcal{P}(^n E, F)$ is compact if maps the unit ball of E into a relatively compact set in F and that P is in $\mathcal{P}_w(^n E, F)$ if maps bounded weak convergent nets into convergent nets. For linear operators both properties, to be compact and to be weakly continuous on bounded sets, produce the same subspace. For n -homogeneous polynomials with $n > 1$, that coincidence is no longer true. Although any polynomial in $\mathcal{P}_w(^n E, F)$ is compact (as it can be derived from results in [9] and [8]), the reverse inclusion fails. This is due to the fact that continuous polynomials are not, in general, weak-to-weak continuous. Then, every scalar-valued continuous polynomial is compact but it is not necessarily weakly continuous on bounded sets, as the standard example $P(x) = \sum_k x_k^2$, for all $x = (x_k)_k \in \ell_2$, shows. With this in mind, our main purpose is to discuss whether $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$. In [16], the first author studied the analogous question when F is the scalar field. We will see that the vector-valued case is not a mere generalization of the scalar-valued case.

The problem of stating if $\mathcal{P}_w(^n E, F)$ is a proper subspace of $\mathcal{P}(^n E, F)$ is nontrivial at all. However, when this is not the situation our question is trivially answered. We refer the reader to [4, 13, 22, 23], where the equality $\mathcal{P}_w(^n E, F) = \mathcal{P}(^n E, F)$ is studied.

As it happens for n -homogeneous polynomials in the scalar-valued case, the value of n for which $\mathcal{P}_w(^n E, F)$ has the chance to be a nontrivial M -ideal in $\mathcal{P}(^n E, F)$ cannot be chosen arbitrarily. Thus, our firsts efforts are focused to discuss this matter. In order to do so, following [24] and [16], we define the essential norm

of a vector-valued polynomial P as the distance from P to the space $\mathcal{P}_w(^nE, F)$. Also we describe the extreme points of the ball of the dual space of $\mathcal{P}_w(^nE, F)$. Then, combining this with properties of the essential norm we obtain the range within we may expect to find an M -structure. When $\mathcal{P}_w(^nE, F)$ is an M -ideal in $\mathcal{P}(^nE, F)$, the essential norm allows us to obtain a Bishop-Phelps type theorem. We use this result to study the existence of farthest points and densely remotal sets. These concepts are related to geometric properties such as the existence of exposed points, the Mazur intersection property and norm attaining functions, see [11, 20]. These results appear in Section 1.

Section 2 is dedicated to give sufficient conditions on E and F so that $\mathcal{P}_w(^nE, F)$ is an M -ideal in $\mathcal{P}(^nE, F)$. The main requirement stays around the concept of shrinking approximations of the identity. When F is an M_∞ -space, without any further assumption on the space E , we prove that $\mathcal{P}_w(^nE, F)$ is a nontrivial M -ideal in $\mathcal{P}(^nE, F)$ for all but one possible value of n in the range of interest. For the remaining value of n , we obtain the result when E satisfies some additional conditions, see Propositions 2.8 and 2.10.

In Section 3, we focus our attention on classical sequence spaces E and F , for $E = \ell_p$ ($1 < p < \infty$) and $F = \ell_q$ or $F = d(w, q)$ a Lorentz sequence space, ($1 < q < \infty$). The questions of whether $\mathcal{K}(\ell_p, \ell_q)$ is an M -ideal in $\mathcal{L}(\ell_p, \ell_q)$ and $\mathcal{K}(\ell_p, d(w, q))$ is an M -ideal in $\mathcal{L}(\ell_p, d(w, q))$ were previously addressed in [24] and [30]. In [16], it was studied when $\mathcal{P}_w(^n\ell_p)$ is an M -ideal in $\mathcal{P}(^n\ell_p)$. We analyze here when $\mathcal{P}_w(^n\ell_p, \ell_q)$ is an M -ideal in $\mathcal{P}(^n\ell_p, \ell_q)$ and when $\mathcal{P}_w(^n\ell_p, d(w, q))$ is an M -ideal in $\mathcal{P}(^n\ell_p, d(w, q))$. Giving conditions on n, p, q and w we solve the problem for all the possible situations.

In the last section we study the property (M) , introduced by Kalton in [26] for Banach spaces, developed later for operators by Kalton and Werner in [27] and finally generalized to the scalar-valued polynomial setting in [16]. Here, we present a natural extension to the vector-valued polynomial setting of the notions mentioned before and establish the connection this property has with our main problem. We apply the results obtained to give examples of M -ideals in vector-valued polynomial spaces defined on Bergman spaces.

Before proceeding, we fix some notation and give basic definitions. Every time we write X, E or F we will be considering Banach spaces over the real or complex field, \mathbb{K} . The closed unit ball of X will be noted by B_X and the unit sphere by S_X . Also, if $x \in X$ and $r > 0$, $B(x, r)$ will stand for the closed ball in X with center at x and radius r . As usual, X^* and X^{**} will be the notations for the dual and bidual of X , respectively. The space of linear bounded operators from E to E will be noted by $\mathcal{L}(E)$ and its subspace of compact mappings will be noted by $\mathcal{K}(E)$.

A function $P: E \rightarrow F$ is said to be an n -homogeneous polynomial if there exists a (unique) symmetric n -linear form $\check{P}: \underbrace{E \times \dots \times E}_n \rightarrow F$ such that

$$P(x) = \check{P}(x, \dots, x),$$

for all $x \in E$. As we have already mentioned, the space of continuous n -homogeneous

polynomials from E to F will be denoted by $\mathcal{P}(^n E, F)$. For scalar-valued mappings we will write $\mathcal{P}(^n E)$ instead of $\mathcal{P}(^n E, \mathbb{K})$ to denote the space of all continuous n -homogeneous polynomials from E to \mathbb{K} . The space $\mathcal{P}(^n E, F)$ endowed with the supremum norm

$$\|P\| = \sup\{\|P(x)\|_F : x \in B_E\},$$

is a Banach space. We may write $\|P(x)\|$ instead of $\|P(x)\|_F$ unless we prefer to emphasize the space where the norm is taken.

Every polynomial $P \in \mathcal{P}(^n E, F)$ has two natural mappings associated: the *linear adjoint* or *transpose* $P^* \in \mathcal{L}(F^*, \mathcal{P}(^n E))$ which is given by

$$(P^*(y^*))(x) = y^*(P(x)), \quad \text{for every } x \in E \text{ and } y^* \in F^*,$$

and the polynomial $\bar{P} \in \mathcal{P}(^n E^{**}, F^{**})$, the canonical extension of P from E to E^{**} obtained by weak-star density, known as the Aron-Berner extension of P [5]. For each $z \in E^{**}$, e_z will refer to the application given by $e_z(P) = \bar{P}(z)$; for $x \in E$, e_x denotes the evaluation map.

Besides the subspace of weakly continuous on bounded sets n -homogeneous polynomials which was already introduced, we will consider the following classes. The first one is the space of n -homogeneous polynomials that are weakly continuous on bounded sets at 0, which consists on those polynomials mapping bounded weakly null nets into null nets. This space will be denoted by $\mathcal{P}_{w0}(^n E, F)$. We also have the subspace formed by polynomials of finite type, which are of the form $\sum_{j=1}^N (x_j^*)^n \cdot y_j$, with $x_j^* \in E^*$, $y_j \in F$ for all $j = 1, \dots, N$ and $N \in \mathbb{N}$. The space of finite type n -homogeneous polynomials will be denoted by $\mathcal{P}_f(^n E, F)$. Its closure (in the supremum norm) is the space of approximable n -homogeneous polynomials which will be noted by $\mathcal{P}_A(^n E, F)$. When F is \mathbb{K} we omit F and write $\mathcal{P}_{w0}(^n E)$, $\mathcal{P}_f(^n E)$ or $\mathcal{P}_A(^n E)$ for instance.

Recall that if E does not contain a subspace isomorphic to ℓ_1 , then, for any Banach space F , $\mathcal{P}_w(^n E, F)$ coincides with the space of weakly sequentially continuous polynomials $\mathcal{P}_{wsc}(^n E, F)$ [8, Proposition 2.12]. The space of n -homogeneous polynomials that are weakly sequentially continuous at 0 will be denoted $\mathcal{P}_{wsc0}(^n E, F)$. As usual, $\mathcal{P}(E, F)$ ($\mathcal{P}_w(E, F)$) stands for the space of all continuous (weakly continuous on bounded sets) polynomials from E to F . We refer to [18, 29] for the necessary background on polynomials on Banach spaces.

Related to the study of M -structures there are two relevant geometric properties that we will use repeatedly. The first one is a well-known characterization, called *the 3-ball property*, given by Alfsen and Effros in [3, Theorem A] to which the main part of their article is dedicated, see also [24, Theorem I.2.2 (iv)]:

Theorem A. *Suppose that J is a closed subspace of X . The following are equivalent:*

- (i) J is an M -ideal.
- (ii) J satisfies the 3-ball property: for every $x_1, x_2, x_3 \in X$ and positive numbers

r_1, r_2, r_3 such that

$$\bigcap_{j=1}^3 B(x_j, r_j) \neq \emptyset \quad \text{and} \quad B(x_j, r_j) \cap J \neq \emptyset, \quad j = 1, 2, 3,$$

it holds that

$$\bigcap_{j=1}^3 B(x_j, r_j + \varepsilon) \cap J \neq \emptyset \quad \text{for all } \varepsilon > 0.$$

(iii) J satisfies the (restricted) 3-ball property: for every $y_1, y_2, y_3 \in B_J$, $x \in B_X$ and $\varepsilon > 0$, there exists $y \in J$ satisfying

$$\|x + y_j - y\| \leq 1 + \varepsilon, \quad j = 1, 2, 3.$$

Note that one of the benefits of having the 3-ball property is that we have a criterium to decide if a closed subspace of a Banach space X is an M -ideal in terms of an intersection of balls in X . Thus, there is no need to appeal to the dual space to determine the existence of an M -structure. The 2-ball property is not sufficient to this end, see [24]. When a closed subspace of X satisfies the 2-ball property we say that we are in presence of a **semi M -ideal** structure.

The second property we referred, provides us with a nice description of the extreme points of the unit ball of X^* in terms of the sets of the extreme points of the unit balls of J^\perp and J^* , if J is an M -ideal in X , see [24, Lemma 1.5]. As usual $\text{Ext}(B_X)$ denotes the set of extreme points of the unit ball of a Banach space X .

Theorem B. *Suppose that J is an M -ideal in X . Then, the extreme points of the unit ball of X^* satisfy*

$$\text{Ext}(B_{X^*}) = \text{Ext}(B_{J^\perp}) \cup \text{Ext}(B_{J^*}).$$

Many authors investigated M -structures on Banach spaces. Hardmand, Werner and Werner summarized the main results on this topic in their monograph [24]. The reader will find out that it is a very clear and well-organized survey on M -ideals. Along this paper, we will recourse to the ideas and results in it.

1. General results

It is natural to begin our research with vector-valued polynomial versions of basic results stated for linear operators in [24, Propositions VI.4.2 and VI.4.3] and for scalar-valued polynomials in [16, Propositions 1.1 and 1.2]. We omit the proofs since they are straightforward.

Proposition 1.1.

- (a) *If $\mathcal{P}_w({}^n E, F)$ is an M -summand in $\mathcal{P}({}^n E, F)$, then $\mathcal{P}_w({}^n E, F) = \mathcal{P}({}^n E, F)$.*
- (b) *If $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$ and $E_1 \subset E$, $F_1 \subset F$ are 1-complemented subspaces, then $\mathcal{P}_w({}^n E_1, F_1)$ is an M -ideal in $\mathcal{P}({}^n E_1, F_1)$.*

- (c) *The class of Banach spaces E and F for which $\mathcal{P}_w(^nE, F)$ is an M -ideal in $\mathcal{P}(^nE, F)$ is closed with respect to the Banach-Mazur distance.*

The knowledge of the extreme points of the unit ball of a Banach space provides a crucial tool in the geometric study of the space. We borrow some ideas of [24] and [16] to examine the extreme points of the unit ball of the dual spaces: $\mathcal{P}(^nE, F)^*$ and $\mathcal{P}_w(^nE, F)^*$.

Note that if J is a subspace of $\mathcal{P}(^nE, F)$ that contains $\mathcal{P}_f(^nE, F)$, then $e_x \otimes y^* \in J^*$ is a norm one element, for all $x \in S_E$ and $y^* \in S_{F^*}$. Indeed, the application $e_x \otimes y^*$ belongs to B_{J^*} and since J contains all finite type n -homogeneous polynomials, it contains the elements of the form $(x^*)^n \cdot y$, for every $x^* \in E^*$ and $y \in F$, thus $\|e_x \otimes y^*\| = 1$.

Proposition 1.2.

- (a) *If J is a subspace of $\mathcal{P}(^nE, F)$ that contains all finite type n -homogeneous polynomials, then*

$$\text{Ext } B_{J^*} \subset \overline{\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*},$$

where w^* designates the topology $\sigma(J^*, J)$.

- (b) *For the particular case $J = \mathcal{P}_w(^nE, F)$ we can be more precise:*

$$\text{Ext } B_{\mathcal{P}_w(^nE, F)^*} \subset \{e_z \otimes y^* : z \in S_{E^{**}}, y^* \in S_{F^*}\}.$$

Proof. (a) Through Hahn-Banach theorem and the comment made above, it easily follows that

$$B_{J^*} = \overline{\Gamma\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*}.$$

Now, by Milman’s theorem [21, Theorem 3.41] we derive the desired inclusion:

$$\text{Ext } B_{J^*} \subset \overline{\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*}.$$

- (b) Suppose that $J = \mathcal{P}_w(^nE, F)$. Let us see that $\overline{\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*} \subset \{e_z \otimes y^* : z \in B_{E^{**}}, y^* \in B_{F^*}\}$. If $\Phi \in \overline{\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*}$, then there exist nets $\{x_\alpha\}_\alpha$ in S_E and $\{y_\alpha^*\}_\alpha$ in S_{F^*} such that $e_{x_\alpha} \otimes y_\alpha^* \xrightarrow{w^*} \Phi$. Without loss of generality, we may assume that $\{x_\alpha\}_\alpha$ is $\sigma(E^{**}, E^*)$ -convergent to an element z in $B_{E^{**}}$ and $\{y_\alpha^*\}_\alpha$ is $\sigma(F^*, F)$ -convergent to an element y^* in B_{F^*} .

Note that for any $P \in \mathcal{P}_w(^nE, F)$, its Aron-Berner extension \overline{P} belongs to $\mathcal{P}(^nE^{**}, F)$ (see for instance [14, Proposition 2.5]) and the compactity of P implies that \overline{P} is w^* -continuous on bounded sets. Then, we have that $y_\alpha^* (\overline{P}(x_\alpha)) \rightarrow y^* (\overline{P}(z))$, for every $P \in \mathcal{P}_w(^nE, F)$. Thus, $e_{x_\alpha} \otimes y_\alpha^* \xrightarrow{w^*} e_z \otimes y^*$ and therefore, $\Phi = e_z \otimes y^*$. When Φ is a norm one element we have that both z and y are elements in the respective unit spheres $S_{E^{**}}$ and S_{F^*} . Now, the result follows. □

In [16], the notion of the essential norm was extended from operators to scalar-valued polynomials and was used to determine that $\mathcal{P}_w(^nE)$ may be a nontrivial M -ideal of $\mathcal{P}(^nE)$ for at most only one value of n . For vector-valued polynomials, also through the essential norm, we obtain a finite range of possible values of n for which $\mathcal{P}_w(^nE, F)$ has the chance to be a nontrivial M -ideal of $\mathcal{P}(^nE, F)$. Recall that the essential norm of a linear operator T is the distance from T to the subspace of compact operators. When $\mathcal{K}(E, F)$ is an M -ideal in $\mathcal{L}(E, F)$, there is an explicit alternative formula to compute this essential norm [24, Proposition VI.4.7]. Now we proceed to discuss the degrees of homogeneity for which our problem might have a nontrivial solution.

Definition 1.3. Let $P \in \mathcal{P}(^nE, F)$ be an n -homogeneous polynomial. The **essential norm** of P is defined by

$$\|P\|_{es} = d(P, \mathcal{P}_w(^nE, F)) = \inf\{\|P - Q\| : Q \in \mathcal{P}_w(^nE, F)\}.$$

In order to obtain a good description of the essential norm, we will make use of the transpose of a polynomial. Note that if $P \in \mathcal{P}(^nE, F)$ and we denote by $L_P : \bigotimes_{\pi_s}^{n,s} E \rightarrow F$ the linearization of P , where π_s is the projective symmetric tensor norm; then P^* is the usual adjoint of L_P .

Lemma 1.4. *If $P \in \mathcal{P}_w(^nE, F)$ then P^* belongs to $\mathcal{L}(F^*, \mathcal{P}_w(^nE))$ and it is w^* -continuous on bounded sets.*

Proof. If $P \in \mathcal{P}_w(^nE, F)$ then P is compact and $P^* \in \mathcal{L}(F^*, \mathcal{P}_w(^nE))$. By [10, Proposition 3.2], P^* is a compact operator. Since $P^* = L_P^*$ it follows that L_P is compact and its adjoint P^* is w^* -continuous. \square

Now we can obtain an alternative formula for the essential norm in the case that there is an M -structure.

Proposition 1.5. *Suppose $\mathcal{P}_w(^nE, F)$ is an M -ideal in $\mathcal{P}(^nE, F)$. Then, for any $P \in \mathcal{P}(^nE, F)$,*

$$\|P\|_{es} = \max\{w(P), w^*(P)\},$$

where

$$w(P) = \sup \left\{ \limsup \|P(x_\alpha)\| : \|x_\alpha\| = 1, x_\alpha \xrightarrow{w} 0 \right\} \quad \text{and}$$

$$w^*(P) = \sup \left\{ \limsup \|P^*(y_\alpha^*)\| : \|y_\alpha^*\| = 1, y_\alpha^* \xrightarrow{w^*} 0 \right\}.$$

Proof. Let $P \in \mathcal{P}(^nE, F)$. For any $Q \in \mathcal{P}_w(^nE, F)$ and for any normalized weak-star null net $\{y_\alpha^*\}_\alpha$, it holds

$$\|P - Q\| = \|P^* - Q^*\| \geq \|(P^* - Q^*)(y_\alpha^*)\| \geq \|P^*(y_\alpha^*)\| - \|Q^*(y_\alpha^*)\|.$$

Since, by Lemma 1.4, $\|Q^*(y_\alpha^*)\| \rightarrow 0$ it follows that $\|P - Q\| \geq \limsup \|P^*(y_\alpha^*)\|$ and thus $\|P\|_{es} \geq w^*(P)$.

The other inequality follows analogously. Thus,

$$\|P\|_{es} \geq \max\{w(P), w^*(P)\}.$$

Now suppose that $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$. Then we have

$$\text{Ext } B_{\mathcal{P}({}^n E, F)^*} = \text{Ext } B_{\mathcal{P}_w({}^n E, F)^\perp} \cup \text{Ext } B_{\mathcal{P}_w({}^n E, F)^*}.$$

The essential norm of P , $\|P\|_{es}$, is the norm of the class of P in the quotient space $\mathcal{P}({}^n E, F)/\mathcal{P}_w({}^n E, F)$ and the dual of this quotient can be isometrically identified with $\mathcal{P}_w({}^n E, F)^\perp$. Then, there exists $\Phi \in \text{Ext } B_{\mathcal{P}_w({}^n E, F)^\perp}$ such that $\Phi(P) = \|P\|_{es}$. So, $\Phi \in \text{Ext } B_{\mathcal{P}({}^n E, F)^*}$ and, by Proposition 1.2(a), $\Phi \in \overline{\{e_x \otimes y^* : x \in S_E, y^* \in S_{F^*}\}}^{w^*}$.

Chose nets $\{x_\alpha\}_\alpha$ in S_E and $\{y_\alpha^*\}_\alpha$ in S_{F^*} such that $e_{x_\alpha} \otimes y_\alpha^* \xrightarrow{w^*} \Phi$, where w^* means the topology $\sigma(\mathcal{P}({}^n E, F)^*, \mathcal{P}({}^n E, F))$. In passing to appropriate subnets, we can suppose that $\{x_\alpha\}_\alpha$ is $\sigma(E^{**}, E^*)$ -convergent to an element z in $B_{E^{**}}$ and $\{y_\alpha^*\}_\alpha$ is $\sigma(F^*, F)$ -convergent to an element y in B_{F^*} .

For any $x^* \in E^*$ and $y \in F$, the polynomial $(x^*)^n \cdot y$ belongs to $\mathcal{P}_w({}^n E, F)$. This gives

$$0 = \Phi((x^*)^n \cdot y) = \lim_\alpha x^*(x_\alpha)^n y_\alpha^*(y) = z(x^*)^n y^*(y).$$

So it should be $z = 0$ or $y^* = 0$. In the first case, $\{x_\alpha\}_\alpha$ is weakly null and

$$\|P\|_{es} = \Phi(P) = \lim_\alpha y_\alpha^*(P(x_\alpha)) \leq \limsup \|P(x_\alpha)\| \leq w(P).$$

In the second case, $\{y_\alpha^*\}_\alpha$ is weak-star null and it follows similarly that $\|P\|_{es} \leq w^*(P)$. □

As in the scalar-valued polynomial case this result enable us to narrow the possible values of n for which $\mathcal{P}_w({}^n E, F)$ could be an M -ideal in $\mathcal{P}({}^n E, F)$. To see this, we extend the definition of critical degree of a Banach space $\text{cd}(E)$ given in [16] to the case of vector-valued polynomials.

If $\mathcal{P}_w(E, F) \neq \mathcal{P}(E, F)$ we define **the critical degree of (E, F)** as:

$$\text{cd}(E, F) = \min\{k \in \mathbb{N}; \mathcal{P}_w({}^k E, F) \neq \mathcal{P}({}^k E, F)\}.$$

Note that if $F = \mathbb{K}$ then $\text{cd}(E) = \text{cd}(E, \mathbb{K})$.

Remark 1.6. Since the same arguments from [13] and [6] used to state Remark 1.8 in [16] also work for vector-valued polynomials we obtain that for Banach spaces E and F if $\mathcal{P}_w(E, F) \neq \mathcal{P}(E, F)$ and $n = \text{cd}(E, F)$,

- $\mathcal{P}_w({}^k E, F) = \mathcal{P}_{w0}({}^k E, F) = \mathcal{P}({}^k E, F)$, for all $k < n$.
- $\mathcal{P}_w({}^n E, F) = \mathcal{P}_{w0}({}^n E, F) \subsetneq \mathcal{P}({}^n E, F)$.
- $\mathcal{P}_w({}^k E, F) \subsetneq \mathcal{P}_{w0}({}^k E, F) \subset \mathcal{P}({}^k E, F)$, for all $k > n$.

Observe that if a scalar-valued polynomial $P \in \mathcal{P}({}^n E)$ is not weakly continuous on bounded sets then, for any $y \in F$, $y \neq 0$, the polynomial $x \mapsto P(x)y$ belongs to $\mathcal{P}({}^n E, F)$ and it is not weakly continuous on bounded sets. This says that, for any Banach space F ,

$$\text{cd}(E, F) \leq \text{cd}(E).$$

Note also that $\text{cd}(E, F)$ could be much smaller than $\text{cd}(E)$. For instance, $\text{cd}(\ell_p, c_0) = 1$ while $\text{cd}(\ell_p)$ is the integer number satisfying $p \leq \text{cd}(\ell_p) < p + 1$.

Example 1.7. Let $E = \ell_p$ and $F = \ell_q$, $1 < p, q < \infty$ or, more generally, $E = \bigoplus_{\ell_p} X_m$ and $F = \bigoplus_{\ell_q} Y_m$, where X_m and Y_m are finite dimensional spaces. From [23] we can derive that the critical degree is the integer number $\text{cd}(E, F)$ satisfying $\frac{p}{q} \leq \text{cd}(E, F) < \frac{p}{q} + 1$.

Lemma 1.8. *Let $P \in \mathcal{P}(^n E, F)$ be a compact polynomial.*

- (a) *If $n < \text{cd}(E)$ then P is weakly continuous on bounded sets.*
- (b) *$w^*(P) = 0$.*

Proof. (a) If $n < \text{cd}(E)$, then every scalar-valued n -homogeneous polynomial on E is weakly continuous on bounded sets. Then, P is weak-to-weak continuous on bounded sets. So, for any bounded net $\{x_\alpha\}_\alpha$ in E such that $x_\alpha \xrightarrow{w} x$, we have $P(x_\alpha) \xrightarrow{w} P(x)$. Being P compact, the bounded net $\{P(x_\alpha)\}_\alpha$ should have a convergent subnet. By a canonical argument we derive that $P(x_\alpha) \rightarrow P(x)$ and thus P is weakly continuous on bounded sets.

(b) This is a consequence of the proof of Lemma 1.4. □

Proposition 1.9. *Every polynomial in $\mathcal{P}(^n E, F)$ which is weakly continuous on bounded sets at 0 and compact is weakly continuous on bounded sets if and only if $n \leq \text{cd}(E)$.*

Proof. If $n > \text{cd}(E)$, there exists a polynomial $p \in \mathcal{P}_{w0}(^n E) \setminus \mathcal{P}_w(^n E)$. Then, for a fixed non null $y \in F$ the polynomial $P(x) = p(x)y$ is weakly continuous on bounded sets at 0 and compact but it is not weakly continuous on bounded sets.

Reciprocally, let $n \leq \text{cd}(E)$ and let $P \in \mathcal{P}(^n E, F)$ be a polynomial weakly continuous on bounded sets at 0 and compact. We know from [10, Proposition 3.4] that, for $0 < k < n$, any derivative $d^k P(x)$ is compact. Thus, by Lemma 1.8(a), we obtain that $d^k P(x)$ is weakly continuous on bounded sets, for all $0 < k < n$. This fact together with the hypothesis of P being weakly continuous on bounded sets at 0 implies that P is weakly continuous on bounded sets. □

The previous results allow us to obtain an upper bound for the numbers n such that $\mathcal{P}_w(^n E, F)$ could be an M -ideal in $\mathcal{P}(^n E, F)$.

Corollary 1.10. *If $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$, then $n \leq \text{cd}(E)$.*

Proof. By Lemma 1.8(b), if $P \in \mathcal{P}(^n E, F)$ is weakly continuous on bounded sets at 0 and compact then $w(P) = w^*(P) = 0$. If, in addition, $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$, Proposition 1.5 states that $\|P\|_{es} = 0$ and so P is weakly continuous on bounded sets. Thus, by Proposition 1.9, it should be $n \leq \text{cd}(E)$. □

Remark 1.11. Clearly, if $n < \text{cd}(E, F)$, $\mathcal{P}_w(^n E, F)$ is a trivial M -ideal in $\mathcal{P}(^n E, F)$. On the other hand, by Corollary 1.10 if $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$ then $n \leq \text{cd}(E)$. Therefore, the problem of whether $\mathcal{P}_w(^n E, F)$ is an M -ideal is worth being studied only for polynomials of degree n , with $\text{cd}(E, F) \leq n \leq \text{cd}(E)$.

The fact that $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$ has some incidence in the set of polynomials whose Aron-Berner extension attains the norm. As we have for scalar-valued polynomials [16, Proposition 1.10], the following version of [24, Proposition VI.4.8] is a Bishop-Phelps type result for vector-valued polynomials. The proof is omitted since it can be obtained as a slight modification of the proof given in [16].

Proposition 1.12. *Let E and F be Banach spaces and suppose that $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.*

- (a) *If $P \in \mathcal{P}({}^n E, F)$ is such that its Aron-Berner extension \bar{P} does not attain its norm at $B_{E^{**}}$, then $\|P\| = \|P\|_{es}$.*
- (b) *The set of polynomials in $\mathcal{P}({}^n E, F)$ whose Aron-Berner extension does not attain the norm is nowhere dense in $\mathcal{P}({}^n E, F)$.*

We finish this section relating norm attaining polynomials with farthest points and remotal sets. The study of the existence of farthest points in a set of a Banach space can be traced to the articles of Asplund [11] and Edelstein [20]. This concept is related to several geometric properties of the space, like the existence of exposed points and the Mazur intersection property.

Perhaps some definitions are in order. Let J be a subspace of a Banach space X . Fix $x \in X$, the **farthest distance** from x to the unit ball of J is given by

$$\rho(x, B_J) = \sup\{\|x - y\| : y \in B_J\}.$$

A point $x \in X$ has a **farthest point** in B_J if there exists $y \in B_J$ such that $\|x - y\| = \rho(x, B_J)$. The set of points in X having farthest points in B_J is denoted by $R(B_J)$. Then we have:

$$R(B_J) = \{x \in X : \exists y \in B_J \text{ such that } \|x - y\| = \rho(x, B_J)\}.$$

It is said that B_J is **densely remotal** in X if $R(B_J)$ is dense in X and it is **almost remotal** in X if $R(B_J)$ contains a dense G_δ set.

In [12], Bandyopadhyay, Lin and Rao studied dense remotality of the ball of $\mathcal{K}(E, F)$ in the space $\mathcal{L}(E, F)$. Adapting some of their ideas and applying the previous proposition, in Corollary 1.17, we obtain a result about almost remotality of $B_{\mathcal{P}_w({}^n E, F)}$ in $\mathcal{P}({}^n E, F)$.

Lemma 1.13. *For any $P \in \mathcal{P}({}^n E, F)$ we have that*

$$\rho(P, B_{\mathcal{P}_w({}^n E, F)}) = \|P\| + 1.$$

Proof. It is clear that $\rho(P, B_{\mathcal{P}_w({}^n E, F)}) \leq \|P\| + 1$, for every $P \in \mathcal{P}({}^n E, F)$ and the equality is obvious for the polynomial $P \equiv 0$. For the reverse inequality, given $P \in \mathcal{P}({}^n E, F)$, $P \not\equiv 0$, and $\varepsilon > 0$, fix $x \in S_E$ and $y^* \in S_{F^*}$ such that $y^*(P(x)) > (1 - \varepsilon)\|P\|$. Now, take $y \in S_F$ and $x^* \in S_{E^*}$ satisfying $y^*(y) > 1 - \varepsilon$ and $x^*(x) = 1$ and consider the polynomial $Q = -(x^*)^n \cdot y \in B_{\mathcal{P}_w({}^n E, F)}$.

Then, we have

$$\begin{aligned} \|P - Q\| &= \|P + (x^*)^n \cdot y\| \geq |y^*(P(x)) + x^*(x)^n y^*(y)| \\ &= y^*(P(x)) + y^*(y) > (1 - \varepsilon)(\|P\| + 1), \end{aligned}$$

for all $\varepsilon > 0$, which proves the lemma. □

The relation between norm attaining linear functions and the set of operators which admit farthest points in the unit ball of the space of compact operators was studied in [12]. To simplify our statements let us introduce the following notations:

$$NA(\mathcal{P}({}^n E, F)) = \{P \in \mathcal{P}({}^n E, F) : P \text{ attains its norm at } B_E\},$$

$$AB - NA(\mathcal{P}({}^n E, F)) = \{P \in \mathcal{P}({}^n E, F) : \bar{P} \text{ attains its norm at } B_{E^{**}}\}.$$

Proposition 1.14. $NA(\mathcal{P}({}^n E, F)) \subset R(B_{\mathcal{P}_w({}^n E, F)})$.

Proof. By the previous lemma, it is plain that the polynomial $P \equiv 0$ belongs to $R(B_{\mathcal{P}_w({}^n E, F)})$. Now, if $P \in NA(\mathcal{P}({}^n E, F))$, $P \neq 0$, there exists $x \in S_E$ such that $\|P(x)\| = \|P\|$. Let $x^* \in S_{E^*}$ satisfying $x^*(x) = 1$.

Consider the polynomial $Q = -(x^*)^n \cdot \frac{P(x)}{\|P\|} \in B_{\mathcal{P}_w({}^n E, F)}$. So Q is a farthest point for P because

$$\|P - Q\| = \left\| P + (x^*)^n \cdot \frac{P(x)}{\|P\|} \right\| \geq \left\| P(x) + \frac{P(x)}{\|P\|} \right\| = \|P\| + 1. \quad \square$$

In [15], Choi and Kim proved that if E has the Radon-Nykodým property, then the set of norm attaining polynomials of $\mathcal{P}({}^n E, F)$ is dense in $\mathcal{P}({}^n E, F)$. As a consequence of this result we obtain:

Corollary 1.15. *If E has the Radon-Nykodým property, then $B_{\mathcal{P}_w({}^n E, F)}$ is densely remotal in $\mathcal{P}({}^n E, F)$.*

When $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$, the set $R(B_{\mathcal{P}_w({}^n E, F)})$ does not only contain the set of norm attaining polynomials but it is also contained in the set of all the polynomials whose Aron-Berner extension is norm attaining.

Proposition 1.16. *If $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$, then $R(B_{\mathcal{P}_w({}^n E, F)}) \subset AB - NA(\mathcal{P}({}^n E, F))$.*

Proof. Let $P \in R(B_{\mathcal{P}_w({}^n E, F)})$. So, there exists $Q \in B_{\mathcal{P}_w({}^n E, F)}$ such that $\|P - Q\| = \|P\| + 1$. Take $\Phi \in \text{Ext } B_{\mathcal{P}({}^n E, F)^*}$ satisfying

$$\Phi(P - Q) = \|P - Q\| = \|P\| + 1.$$

Being $\mathcal{P}_w({}^n E, F)$ an M -ideal in $\mathcal{P}({}^n E, F)$, we should have that

$$\Phi \in \text{Ext } B_{\mathcal{P}_w({}^n E, F)^*} \quad \text{or} \quad \Phi \in \text{Ext } B_{\mathcal{P}_w({}^n E, F)^\perp}.$$

If $\Phi \in \text{Ext } B_{\mathcal{P}_w({}^n E, F)^\perp}$, we obtain that $\Phi(P - Q) = \Phi(P)$ and so $\Phi(P) = \|P\| + 1$, which is not possible. Hence, it should be $\Phi \in \text{Ext } B_{\mathcal{P}_w({}^n E, F)^*}$ and, by Proposition 1.2(b), $\Phi = e_z \otimes y^*$, for certain $z \in S_{E^{**}}$ and $y^* \in S_{F^*}$. Therefore,

$$\|P\| + 1 = \Phi(P - Q) = y^*(\bar{P}(z)) - y^*(\bar{Q}(z)) \leq \|\bar{P}\| + \|\bar{Q}\| = \|P\| + 1.$$

It follows that $y^*(\bar{P}(z)) = \|\bar{P}\|$ and so $\|\bar{P}(z)\| = \|\bar{P}\|$, meaning that $P \in AB - NA(\mathcal{P}({}^n E, F))$. □

As a consequence of Propositions 1.12, 1.14 and 1.16, we obtain:

Corollary 1.17. *If E is reflexive and $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$, then*

$$R(B_{\mathcal{P}_w({}^n E, F)}) = NA(\mathcal{P}({}^n E, F)),$$

and thus, $\mathcal{P}({}^n E, F) \setminus R(B_{\mathcal{P}_w({}^n E, F)})$ is nowhere dense. This implies that $B_{\mathcal{P}_w({}^n E, F)}$ is almost remotal in $\mathcal{P}({}^n E, F)$.

2. Sufficient conditions

In this section we present several kind of sufficient conditions which enable us to ensure that $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$. All of them involve bounded nets of compact operators on E . The following lemma and proposition are the vector-valued versions of [16, Lemma 2.1 and Proposition 2.2], the proofs of which are analogous to those in [16].

Lemma 2.1. *Let E and F be Banach spaces and suppose that there exists a bounded net $\{S_\alpha\}_\alpha$ of linear operators from E to E satisfying $S_\alpha^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$. Then, for all $P \in \mathcal{P}_w({}^n E, F)$, we have that $\|P - P \circ S_\alpha\| \rightarrow 0$.*

Proposition 2.2. *Let E and F be Banach spaces and let $n = \text{cd}(E, F)$. Suppose that there exists a bounded net $\{K_\alpha\}_\alpha$ of compact operators from E to E satisfying the following two conditions:*

- $K_\alpha^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$.
- For all $\varepsilon > 0$ and all α_0 there exists $\alpha > \alpha_0$ such that for every $x \in E$,

$$\|K_\alpha(x)\|^n + \|x - K_\alpha(x)\|^n \leq (1 + \varepsilon)\|x\|^n.$$

Then, $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.

Remark 2.3. A Banach space E is an (M_p) -space ($1 \leq p \leq \infty$) if $\mathcal{K}(E \oplus_p E)$ is an M -ideal in $\mathcal{L}(E \oplus_p E)$. This concept was introduced by Oja and Werner in [31]. By [24, Theorem VI.5.3], if E is an (M_p) -space with $p \leq n$, then there exists a bounded net $\{K_\alpha\}_\alpha$ of compact operators from E to E satisfying both conditions of Proposition 2.2.

Recall that a Banach space E has a finite dimensional decomposition $\{E_j\}_j$ if each E_j is a finite dimensional subspace of E and every $x \in E$ has a unique representation of the form

$$x = \sum_{j=1}^{\infty} x_j, \quad \text{with } x_j \in E_j, \text{ for every } j.$$

Associated to the decomposition there is a bounded sequence of projections $\{\pi_m\}_m$, given by $\pi_m(\sum_{j=1}^{\infty} x_j) = \sum_{j=1}^m x_j$. The decomposition is called shrinking if $\pi_m^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$.

It is clear that in this case $\{\pi_m\}_m$ is a bounded sequence of compact operators that satisfies the first item of the previous proposition. Thus, for spaces with shrinking finite dimensional decompositions we state the following simpler version of Proposition 2.2.

Corollary 2.4. *Let E and F be Banach spaces and let $n = \text{cd}(E, F)$. Suppose that E has a shrinking finite dimensional decomposition with associate projections $\{\pi_m\}_m$ such that:*

- For all $\varepsilon > 0$ and all $m_0 \in \mathbb{N}$ there exists $m > m_0$ such that for every $x \in E$,

$$\|\pi_m(x)\|^n + \|x - \pi_m(x)\|^n \leq (1 + \varepsilon)\|x\|^n.$$

Then, $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.

Note that the hypothesis of Corollary 2.4 are fulfilled for the classical spaces $E = \ell_p$ and $F = \ell_q$, $1 < p, q < \infty$, in the case of $n = \text{cd}(E, F) \geq p$. In this situation $\mathcal{P}_w({}^n \ell_p, \ell_q)$ is an M -ideal in $\mathcal{P}({}^n \ell_p, \ell_q)$.

The conditions in the following theorem were inspired by those of [24, Lemma VI.6.7]. They concern bounded nets of compact operators both in E and in F .

Theorem 2.5. *Let E and F be Banach spaces and suppose that there exist bounded nets of compact operators $\{K_\alpha\}_\alpha \subset \mathcal{K}(E)$ and $\{L_\beta\}_\beta \subset \mathcal{K}(F)$ and numbers $1 < p, q < \infty$ such that:*

- $K_\alpha^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$ and $L_\beta(y) \rightarrow y$, for all $y \in F$.
- For all $\varepsilon > 0$ and all α_0 there exists $\alpha > \alpha_0$ such that for every $x \in E$,

$$\|K_\alpha(x)\|^p + \|x - K_\alpha(x)\|^p \leq (1 + \varepsilon)^p \|x\|^p.$$

- For all $\varepsilon > 0$ and all β_0 there exists $\beta > \beta_0$ such that for every $y_1, y_2 \in F$,

$$\|L_\beta(y_1) + (Id - L_\beta)(y_2)\|^q \leq (1 + \varepsilon)^q (\|y_1\|^q + \|y_2\|^q).$$

Suppose also that $n = \text{cd}(E, F)$ satisfies that $p \leq nq$ and $n < \text{cd}(E)$. Then, $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.

Proof. We prove that the 3-ball property holds. Let $P_1, P_2, P_3 \in B_{\mathcal{P}_w({}^n E, F)}$, $Q \in B_{\mathcal{P}({}^n E, F)}$ and $\varepsilon > 0$. Define $P = Q - (Id - L_\beta)Q(Id - K_\alpha)$. We want to show that P is weakly continuous on bounded sets and $\|Q + P_j - P\| \leq 1 + \varepsilon$, for $j = 1, 2, 3$, for some convenient choice of α and β .

To see that P is weakly continuous on bounded sets, we write $P = Q - Q(Id - K_\alpha) + L_\beta Q(Id - K_\alpha)$. The proof of [16, Proposition 2.2] shows that $Q - Q(Id - K_\alpha)$ is weakly continuous on bounded sets at 0 and since $n = \text{cd}(E, F)$, we have that $Q - Q(Id - K_\alpha)$ belongs to $\mathcal{P}_w({}^n E, F)$. Also, as $L_\beta Q(Id - K_\alpha)$ is a compact polynomial and $n < \text{cd}(E)$, Lemma 1.8(a) says that it is in $\mathcal{P}_w({}^n E, F)$.

Now, to show the $(1 + \varepsilon)$ -bound, consider the inequality

$$\|Q + P_j - P\| \leq \|Q + L_\beta P_j K_\alpha - P\| + \|P_j - L_\beta P_j K_\alpha\|.$$

On the one hand, we have:

$$\begin{aligned} \|P_j - L_\beta P_j K_\alpha\| &\leq \|P_j - P_j K_\alpha\| + \|P_j K_\alpha - L_\beta P_j K_\alpha\| \\ &\leq \|P_j - P_j K_\alpha\| + \|P_j - L_\beta P_j\| \|K_\alpha\|^n. \end{aligned}$$

By Lemma 2.1, $\|P_j - P_j K_\alpha\| \rightarrow 0$ with α . Also, since L_β approximates the identity on compact sets and the P_j 's are compact polynomials, we have that $\|P_j - L_\beta P_j\| \|K_\alpha\|^n \rightarrow 0$ with β , for all α .

Furthermore, we can find α and β such that:

$$\begin{aligned} \|Q + L_\beta P_j K_\alpha - P\| &= \sup_{x \in B_E} \|(Id - L_\beta)Q(Id - K_\alpha)(x) + L_\beta P_j K_\alpha(x)\| \\ &\leq \sup_{x \in B_E} (1 + \varepsilon) (\|Q(Id - K_\alpha)(x)\|^q + \|P_j K_\alpha(x)\|^q)^{\frac{1}{q}} \\ &\leq (1 + \varepsilon) \sup_{x \in B_E} (\|(Id - K_\alpha)(x)\|^{nq} + \|K_\alpha(x)\|^{nq})^{\frac{1}{q}} \\ &\leq (1 + \varepsilon) \sup_{x \in B_E} (\|(Id - K_\alpha)(x)\|^p + \|K_\alpha(x)\|^p)^{\frac{n}{p}} \\ &\leq (1 + \varepsilon)(1 + \varepsilon)^n = (1 + \varepsilon)^{n+1}, \end{aligned}$$

and the result follows. \square

Remark 2.6. If E is an (M_p) -space and F is an (M_q) -space the conditions about the nets of compact operators of the previous theorem are fulfilled.

Proposition 2.7. Let $E = \bigoplus_{\ell_p} X_m$ and $F = \bigoplus_{\ell_q} Y_m$, with X_m and Y_m finite dimensional spaces and $1 < p, q < \infty$. Then, for $n = \text{cd}(E, F)$, $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.

Proof. As we note in Example 1.7, $\text{cd}(E, F)$ is the integer such that $\frac{p}{q} \leq \text{cd}(E, F) < \frac{p}{q} + 1$. Also we know that $\text{cd}(E)$ is the integer satisfying $p \leq \text{cd}(E) < p + 1$. Thus, the result follows from Corollary 2.4 if $\text{cd}(E, F) \geq p$ and from Theorem 2.5 in the case of $\text{cd}(E, F) < p$. \square

In all the previous results (Proposition 2.2, Corollary 2.4 and Theorem 2.5) the M -structure is obtained only in the case $n = \text{cd}(E, F)$. Let us show now some positive results for values of n greater than $\text{cd}(E, F)$.

Proposition 2.8. Let E be a Banach space and F be an (M_∞) -space. If $n < \text{cd}(E)$, then $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.

Proof. Being F an (M_∞) -space, by [24, Theorem VI.5.3], there exists a net $\{L_\beta\}_\beta$ contained in the unit ball of $\mathcal{K}(F)$ satisfying $L_\beta(y) \rightarrow y$ for all $y \in F$ such that for any $\varepsilon > 0$, there exists β_0 with

$$\|L_\beta(y_1) + (Id - L_\beta)(y_2)\| \leq \left(1 + \frac{\varepsilon}{2}\right) \max\{\|y_1\|, \|y_2\|\}, \quad (1)$$

for all $\beta \geq \beta_0$ and for any $y_1, y_2 \in F$. Let $P_1, P_2, P_3 \in B_{\mathcal{P}_w({}^n E, F)}$ and $Q \in B_{\mathcal{P}({}^n E, F)}$, we show that with $P = L_\beta Q$, choosing β properly, the 3-ball property is satisfied.

First, note that by Lemma 1.8(a), P is weakly continuous on bounded sets. Also, $\|Q + P_j - P\| \leq \|Q + L_\beta P_j - P\| + \|P_j - L_\beta P_j\|$. Reasoning as in Theorem 2.5, we

have that $\|P_j - L_\beta P_j\| < \frac{\varepsilon}{2}$ for β large enough. Now, from (1) we obtain

$$\|Q + L_\beta P_j - P\| = \|(Id - L_\beta)Q + L_\beta P_j\| \leq \left(1 + \frac{\varepsilon}{2}\right) \max\{\|Q\|, \|P_j\|\} = \left(1 + \frac{\varepsilon}{2}\right),$$

and the result follows. □

Remark 2.9. Let E be a Banach space such that $\text{cd}(E) > 2$ and let F be an infinite dimensional (M_∞) -space. Then, for any degree n , with $1 \leq n < \text{cd}(E)$, $\mathcal{P}_w(^n E, F)$ is a nontrivial M -ideal in $\mathcal{P}(^n E, F)$. This is a simple consequence of the above proposition and the fact that $\text{cd}(E, F) = 1$.

The next proposition somehow complements Proposition 2.8. It states that if F is an (M_∞) -space, with an additional hypothesis on E , then $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$ also in the case $n = \text{cd}(E)$.

Proposition 2.10. *Let F be an (M_∞) -space and let E be a Banach space. If $n = \text{cd}(E)$ and there exists a bounded net of compact operators $\{K_\alpha\}_\alpha \subset \mathcal{K}(E)$ satisfying both conditions:*

- $K_\alpha^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$.
- For all $\varepsilon > 0$ and all α_0 there exists $\alpha > \alpha_0$ such that for every $x \in E$,

$$\|K_\alpha(x)\|^n + \|x - K_\alpha(x)\|^n \leq (1 + \varepsilon)\|x\|^n,$$

then $\mathcal{P}_w(^n E, F)$ is an M -ideal in $\mathcal{P}(^n E, F)$.

Proof. Let $P_1, P_2, P_3 \in B_{\mathcal{P}_w(^n E, F)}$, $Q \in B_{\mathcal{P}(^n E, F)}$ and $\varepsilon > 0$. We will find $P \in \mathcal{P}_w(^n E, F)$ such that the 3-ball property is satisfied. Reasoning as in Theorem 2.5, we find α and β so that $\|P_j - L_\beta P_j K_\alpha\| < \frac{\varepsilon}{2}$, where $\{L_\beta\}_\beta$ is a net in $\mathcal{K}(F)$, associated to the (M_∞) -space F . Moreover, α and β may be chosen to satisfy at the same time $\|K_\alpha(x)\|^n + \|x - K_\alpha(x)\|^n \leq (1 + \tilde{\varepsilon})\|x\|^n$ and $\|L_\beta(y_1) + (Id - L_\beta)(y_2)\| \leq (1 + \tilde{\varepsilon}) \max\{\|y_1\|, \|y_2\|\}$ for all $y_1, y_2 \in F$, where $\tilde{\varepsilon}$ is such that $(1 + \tilde{\varepsilon})^2 \leq 1 + \frac{\varepsilon}{2}$.

Let P be the polynomial $P = L_\beta(Q - Q(Id - K_\alpha))$. As in the proof of [16, Proposition 2.2], we can see that P is weakly continuous on bounded sets at 0. Since $n = \text{cd}(E)$ and P is compact, we may appeal to Proposition 1.9 to derive that P is weakly continuous on bounded sets.

Also we have,

$$\begin{aligned} \|Q + P_j - P\| &\leq \|Q + L_\beta P_j K_\alpha - P\| + \|P_j - L_\beta P_j K_\alpha\| \\ &\leq \|(Id - L_\beta)Q + L_\beta(P_j K_\alpha + Q(Id - K_\alpha))\| + \frac{\varepsilon}{2} \\ &\leq (1 + \tilde{\varepsilon}) \sup_{x \in B_E} \max\{\|Q(x)\|, \|P_j K_\alpha(x) + Q(x - K_\alpha(x))\|\} + \frac{\varepsilon}{2}. \end{aligned}$$

Now, the hypothesis on E gives us

$$\|P_j K_\alpha(x) + Q(x - K_\alpha(x))\| \leq \|K_\alpha(x)\|^n + \|x - K_\alpha(x)\|^n \leq (1 + \tilde{\varepsilon}),$$

for all $x \in B_E$, and the result follows. □

Example 2.11. Let $E = \ell_p$, with $1 < p < \infty$, and let F be an (M_∞) -space. As a consequence of the previous propositions, since $p \leq \text{cd}(\ell_p)$, $\mathcal{P}_w({}^n\ell_p, F)$ is an M -ideal in $\mathcal{P}({}^n\ell_p, F)$ for all $1 \leq n \leq \text{cd}(\ell_p)$.

3. Polynomials between classical sequence spaces

This section is devoted to study whether $\mathcal{P}_w({}^nE, F)$ is an M -ideal in $\mathcal{P}({}^nE, F)$, for all the values of n between $\text{cd}(E, F)$ and $\text{cd}(E)$, in the cases $E = \ell_p$ and $F = \ell_q$ or F the Lorentz sequence space $F = d(w, q)$, $1 < p, q < \infty$. Recall that given a non increasing sequence $w = (w_j)_j$ of positive real numbers satisfying $w \in c_0 \setminus \ell_1$, the Lorentz sequence space $d(w, q)$ is the space of all sequences $x = (x_j)_j \subset \mathbb{K}$, such that

$$\sup_{\sigma} \sum_{j=1}^{\infty} w_j |x_{\sigma(j)}|^q < \infty,$$

(where σ varies on the set of permutations of \mathbb{N}) endowed with the norm $\|x\|_{d(w,q)} = \sup_{\sigma} \left(\sum_{j=1}^{\infty} w_j |x_{\sigma(j)}|^q \right)^{\frac{1}{q}}$. We will consider weights $w = (w_j)_j$ so that $w_1 = 1$, which implies that the canonical vectors of $d(w, q)$ form a basis of norm 1 elements.

We begin our study with a result about polynomials from a general Banach space E to a Banach space F having a finite dimensional decomposition (FDD) $\{F_n\}_n$. As usual, $\{\pi_m\}_m$ denotes the sequence of projections associated to the decomposition; that is $\pi_m(y) = \sum_{j=1}^m y_j$ for all $y = \sum_{j=1}^{\infty} y_j$, with $y_j \in F_j$. Also, we denote by $\pi^m = Id - \pi_m$. When the FDD is unconditional with unconditional constant 1, we have that $\|\pi^m\| \leq 1$ and $\|\pi_m + \pi^k\| \leq 1$, for all $k \geq m$. In the sequel, we will use, without further mentioning, that for any Banach space E and any $Q \in \mathcal{P}_w({}^nE, F)$, $\|\pi_m Q - Q\| \rightarrow 0$, or equivalently, $\|\pi^m Q\| \rightarrow 0$, both claims can be derived from the fact that Q is compact.

The following proposition gives conditions under which, if F is a Banach space with 1-unconditional FDD, $\mathcal{P}_w({}^nE, F)$ is not a semi M -ideal in $\mathcal{P}({}^nE, F)$. This is a polynomial generalization of [30, Proposition 2] and our proof is modeled on the proof given in that article. From this, it is obviously inferred that $\mathcal{P}_w({}^nE, F)$ is not an M -ideal in $\mathcal{P}({}^nE, F)$.

Proposition 3.1. *Let E and F be Banach spaces such that F has an unconditional FDD with unconditional constant equal to 1 and associated projections $\{\pi_m\}_m$. Suppose that there exist polynomials $P \in \mathcal{P}({}^nE, F)$ and $Q \in \mathcal{P}_w({}^nE, F)$ and numbers $\delta > 0$ and $m_0 \in \mathbb{N}$ such that:*

- $0 < \|Q\| \leq \|P\| < \delta$,
- $\|\pi^m P + Q\| \geq \delta$, for all $m \geq m_0$.

Then, $\mathcal{P}_w({}^nE, F)$ is not a semi M -ideal in $\mathcal{P}({}^nE, F)$.

Proof. Fix $\varepsilon > 0$ so that $\varepsilon < \frac{\delta - \|P\|}{2}$. Since $\|\pi^m Q\| \rightarrow 0$, we may assume that $\|\pi^m Q\| < \frac{\varepsilon}{3}$, for all $m \geq m_0$. Now, fix $m \geq m_0$ and consider the following two closed balls of radius $\|P\|$: $B_1 = B(\pi^m P + Q, \|P\|)$ and $B_2 = B(\pi^m P - Q, \|P\|)$. Note that $\pi^m P \in B_1 \cap B_2$, $Q \in B_1 \cap \mathcal{P}_w({}^nE, F)$ and $-Q \in B_2 \cap \mathcal{P}_w({}^nE, F)$.

If $\mathcal{P}_w({}^n E, F)$ is a semi M -ideal in $\mathcal{P}({}^n E, F)$, then for any $r > \|P\|$, the intersection $B(\pi^m P + Q, r) \cap B(\pi^m P - Q, r) \cap \mathcal{P}_w({}^n E, F)$ is non void. Take $r = \frac{\|P\| + \delta}{2} - \varepsilon$ and suppose that there exists $R \in B(\pi^m P + Q, r) \cap B(\pi^m P - Q, r) \cap \mathcal{P}_w({}^n E, F)$. Since $\|\pi^k R\| \rightarrow 0$, we may choose $k \geq m$ such that $\|\pi^k R\| < \frac{\varepsilon}{3}$. To get a contradiction we estimate $\|\pi^k P + Q\|$. Note that

$$2\|\pi^k P + Q\| \leq \|\pi^k P + \pi_m Q - \pi_m R\| + \|\pi^k P + \pi_m Q + \pi_m R\| + 2\|\pi^m Q\|. \tag{2}$$

From the equality $(\pi_m + \pi^k)(\pi^m P + Q - R) = \pi^k P + \pi_m Q - \pi_m R + \pi^k Q - \pi^k R$, we obtain:

$$\|\pi^k P + \pi_m Q - \pi_m R\| \leq \|\pi_m + \pi^k\| \|\pi^m P + Q - R\| + \|\pi^k Q\| + \|\pi^k R\| < r + \frac{2\varepsilon}{3}.$$

Also, we have that $\|\pi^k P + \pi_m Q + \pi_m R\| = \|\pi^k P - \pi_m Q - \pi_m R\|$, since F has 1-unconditional finite dimensional decomposition. Proceeding as before, we obtain:

$$\|\pi^k P - \pi_m Q - \pi_m R\| \leq \|\pi_m + \pi^k\| \|\pi^m P - Q - R\| + \frac{2\varepsilon}{3} < r + \frac{2\varepsilon}{3}.$$

Finally, using (2), we have that

$$2\delta \leq 2\|\pi^k P + Q\| < 2r + 2\varepsilon = \|P\| + \delta < 2\delta.$$

Thus, we conclude that $\mathcal{P}_w({}^n E, F)$ is not a semi M -ideal in $\mathcal{P}({}^n E, F)$. □

Now we can complete the case $E = \ell_p$ and $F = \ell_q$.

Theorem 3.2. *Let $n = \text{cd}(\ell_p, \ell_q)$.*

- (a) $\mathcal{P}_w({}^n \ell_p, \ell_q)$ is an M -ideal in $\mathcal{P}({}^n \ell_p, \ell_q)$.
- (b) $\mathcal{P}_w({}^k \ell_p, \ell_q)$ is not a semi M -ideal in $\mathcal{P}({}^k \ell_p, \ell_q)$, for all $k > n$.

Proof. Statement (a) follows from Proposition 2.7. To prove statement (b) take $k > n$. We will construct polynomials $P \in \mathcal{P}({}^k \ell_p, \ell_q)$ and $Q \in \mathcal{P}_w({}^k \ell_p, \ell_q)$ satisfying: $\|P\| = \|Q\|$ and $\|\pi^m P + Q\| \geq \delta > \|P\|$, for some $\delta > 0$, where $\{\pi_m\}_m$ is the sequence of projections associated to the canonical basis of ℓ_q and $\pi^m = Id - \pi_m$, for all $m \in \mathbb{N}$.

We have that $k - 1 \geq \text{cd}(\ell_p, \ell_q) \geq \frac{p}{q}$, as shown in Example 1.7, so we may define the continuous k -homogeneous polynomial $P(x) = e_1^*(x)(x_j^{k-1})_{j \geq 2}$. To compute the norm of P , we look, for each $x \in \ell_p$, at the inequality

$$\|P(x)\|_{\ell_q} = |x_1| \left(\sum_{j=2}^{\infty} |x_j|^{(k-1)q} \right)^{\frac{1}{q}} \leq |x_1| \left(\sum_{j=2}^{\infty} |x_j|^p \right)^{\frac{k-1}{p}}.$$

Then, $\|P\| \leq \max\{ab^{k-1} : a^p + b^p = 1, a, b \geq 0\} = \left[\frac{1}{k}(1 - \frac{1}{k})^{k-1}\right]^{\frac{1}{p}}$. Now, considering

$$\tilde{x} = \left(\frac{1}{k}\right)^{\frac{1}{p}} e_1 + \left(1 - \frac{1}{k}\right)^{\frac{1}{p}} e_2,$$

we obtain a norm one element where P attains the bound $[\frac{1}{k}(1 - \frac{1}{k})^{k-1}]^{\frac{1}{p}}$.

Let $Q \in \mathcal{P}_w(k\ell_p, \ell_q)$ be the polynomial $Q(x) = \|P\|e_1^*(x)^k e_1$. It is clear that $\|P\| = \|Q\|$. Take $m \geq 1$, and $\tilde{x} = (\frac{1}{k})^{\frac{1}{p}}e_1 + (1 - \frac{1}{k})^{\frac{1}{p}}e_{m+2}$, then $\|\tilde{x}\|_{\ell_p} = 1$ and

$$\begin{aligned} \|\pi^m P + Q\| &\geq \|(\pi^m P + Q)(\tilde{x})\|_{\ell_q} \\ &= \left\| \left(\frac{1}{k}\right)^{\frac{1}{p}} \left(1 - \frac{1}{k}\right)^{\frac{k-1}{p}} e_{m+1} + \|P\| \left(\frac{1}{k}\right)^{\frac{k}{p}} e_1 \right\|_{\ell_q} = \|P\| \left(1 + \left(\frac{1}{k}\right)^{\frac{kq}{p}}\right)^{\frac{1}{q}}. \end{aligned}$$

Then, with $\delta = \|P\| \left(1 + \left(\frac{1}{k}\right)^{\frac{kq}{p}}\right)^{\frac{1}{q}} > \|P\|$, which is independent of m , we obtain the inequality we were looking for. And the theorem is proved. \square

Now we focus our attention on spaces of polynomials from ℓ_p to $d(w, q)$, $1 < p, q < \infty$. We study whether $\mathcal{P}_w(k\ell_p, d(w, q))$ is an M -ideal in $\mathcal{P}(k\ell_p, d(w, q))$ for $k \geq \text{cd}(\ell_p, d(w, q))$. To this end we extend to the vector-valued case a couple of results of [17] about polynomials from spaces with finite dimensional decompositions.

Lemma 3.3. *Let E be a Banach space which has an unconditional FDD with associated projections $\{\pi_m\}_m$. For any fixed subsequence $\{m_j\}_j$ of \mathbb{N} , let $\sigma_j = \pi_{m_j} - \pi_{m_{j-1}}$, for all j . Given $P \in \mathcal{P}(^n E, F)$, the application*

$$\tilde{P}(x) = \sum_{j=1}^{\infty} P(\sigma_j(x)), \quad \text{for all } x \in E,$$

defines a continuous n -homogeneous polynomial from E to F .

Proof. We first show that the series $\sum_{j=1}^{\infty} P(\sigma_j(x))$ is convergent for every $x \in E$. Indeed, by [17, Proposition 1.3], there exists $C > 0$ such that

$$\begin{aligned} \left\| \sum_{j=N}^M P(\sigma_j(x)) \right\| &\leq \sup_{y^* \in B_{F^*}} \sum_{j=N}^M |y^* \circ P(\sigma_j(x))| \\ &= \sup_{y^* \in B_{F^*}} \sum_{j=N}^M |y^* \circ P(\sigma_j(\pi_{m_M}(x) - \pi_{m_{N-1}}(x)))| \\ &\leq C \|P\| \|\pi_{m_M}(x) - \pi_{m_{N-1}}(x)\|^n, \end{aligned}$$

which converges to 0 with M and N . Then, $\tilde{P}(x)$ is well defined and $\|\tilde{P}\| \leq C\|P\|$. \square

Recall that whenever a Banach space E has a shrinking FDD, by [8], $\mathcal{P}_w(^n E, F) = \mathcal{P}_{wsc}(^n E, F)$. This allows us to work with sequences instead of nets.

Proposition 3.4. *Let E be a Banach space with an unconditional FDD and let F be a Banach space. For any $n \in \mathbb{N}$, the following are equivalent:*

- (i) $\mathcal{P}(^n E, F) = \mathcal{P}_{wsc}(^n E, F)$.

(ii) $\mathcal{P}({}^n E, F) = \mathcal{P}_{wsc0}({}^n E, F)$.

Proof. By means of the previous lemma, the scalar valued result given in [17, Corollary 1.7] (see also [13]) can be easily modified to obtain this vector valued version. □

In [30], Eve Oja studies when $\mathcal{K}(\ell_p, d(w, q))$ is an M -ideal in $\mathcal{L}(\ell_p, d(w, q))$. In Proposition 1 of that article, she establishes a criterium to ensure that every continuous linear operator is compact. A polynomial version of this result can be stated as follows.

Proposition 3.5. *Let $\{e_j\}_j$ and $\{f_j\}_j$ be sequences in Banach spaces E and F , respectively, satisfying:*

- *For any semi-normalized weakly null sequence $\{x_m\}_m \subset E$, there exists a subsequence $\{x_{m_j}\}_j$ and an operator $T \in \mathcal{L}(E)$ such that $T(e_j) = x_{m_j}$, for all j .*
- *For any semi-normalized weakly null sequence $\{y_m\}_m \subset F$, there exists a subsequence $\{y_{m_j}\}_j$ and an operator $S \in \mathcal{L}(F)$ such that $S(y_{m_j}) = f_j$, for all j .*
- *For any subsequence $\{e_{j_l}\}_l$ of $\{e_j\}_j$, there exists an operator $R \in \mathcal{L}(E)$ such that $R(e_l) = e_{j_l}$, for all l .*

Take $n < \text{cd}(E)$ and suppose that it does not exist a polynomial $P \in \mathcal{P}({}^n E, F)$ such that $P(e_j) = f_j$, for every j . Then, $\mathcal{P}({}^n E, F) = \mathcal{P}_{wsc0}({}^n E, F)$.

Proof. Suppose there exists $P \in \mathcal{P}({}^n E, F)$ which is not in $\mathcal{P}_{wsc0}({}^n E, F)$. Then, there exists a weakly null sequence $(x_m)_m$ such that $\|P(x_m)\| > \varepsilon$, for some $\varepsilon > 0$ and all m . As $n < \text{cd}(E)$, $(P(x_m))_m$ is weakly null. Now, we may find a subsequence $(x_{m_j})_j$ and operators $R, T \in \mathcal{L}(E)$ and $S \in \mathcal{L}(F)$ satisfying:

$$e_j \xrightarrow{T \circ R} x_{m_j} \xrightarrow{P} P(x_{m_j}) \xrightarrow{S} f_j,$$

which is a contradiction since $S \circ P \circ T \circ R$ belongs to $\mathcal{P}({}^n E, F)$. □

Remark 3.6. If the Banach space E has an unconditional basis $\{e_j\}_j$ with coordinate functionals $\{e_j^*\}_j$ and $\{f_j\}_j$ is a sequence in the Banach space F , we derive from Lemma 3.3 that the existence of a polynomial $P \in \mathcal{P}({}^n E, F)$ such that $P(e_j) = f_j$, for all j , is equivalent to the existence of the polynomial $\tilde{P} \in \mathcal{P}({}^n E, F)$ given by

$$\tilde{P}(x) = \sum_{j=1}^{\infty} (e_j^*(x))^n f_j, \quad \text{for all } x \in E.$$

When E and F are Banach sequence spaces with canonical bases $\{e_j\}_j$ and $\{f_j\}_j$ respectively, we write the polynomial above as $\tilde{P}(x) = (x_j^n)_j$.

Let $1 < p, q < \infty$. To study whether $\mathcal{P}_w({}^n \ell_p, d(w, q))$ is an M -ideal in $\mathcal{P}({}^n \ell_p, d(w, q))$ for $n \geq \text{cd}(\ell_p, d(w, q))$, we need first to establish the value of the critical degree,

$\text{cd}(\ell_p, d(w, q))$. To this end and in view of the previous remark and proposition, the point is to determine the values of n , p and q such that the polynomial $x \mapsto (x_j^n)_j$, from ℓ_p to $d(w, q)$, is well defined. For $1 < r < \infty$ we use the standard notation $s = r^*$ to denote the conjugate number of r : $\frac{1}{r} + \frac{1}{s} = 1$.

Proposition 3.7. *The polynomial $P(x) = (x_j^n)_j$ belongs to $\mathcal{P}(^n\ell_p, d(w, q))$ if and only if one of the following two conditions holds:*

(a) $n \geq \frac{p}{q}$. In this case, $\|P\| = 1$.

(b) $n < \frac{p}{q}$ and $w \in \ell_s$, for $s = (\frac{p}{nq})^*$. In this case, $\|P\| = \|w\|_{\ell_s}^{\frac{1}{q}}$.

Proof. Let $(e_j)_j$ and $(f_j)_j$ be the canonical bases of ℓ_p and $d(w, q)$, respectively. Suppose that $n \geq \frac{p}{q}$, as $\|w\|_\infty = 1$, we have

$$\|P(x)\|_{d(w,q)} = \sup_{\sigma} \left(\sum_{j=1}^{\infty} w_j |x_{\sigma(j)}|^{nq} \right)^{\frac{1}{q}} \leq \|x\|_{\ell_p}^n.$$

Then, P is a well defined polynomial with norm less than or equal to 1. Also, $P(e_j) = f_j$ implies $\|P\| = 1$.

Now, suppose that $n < \frac{p}{q}$ and $w \in \ell_s$, with $s = (\frac{p}{nq})^*$. Put $W = \|w\|_{\ell_s}$, by Hölder inequality, we have

$$\|P(x)\|_{d(w,q)} = \sup_{\sigma} \left(\sum_{j=1}^{\infty} w_j |x_{\sigma(j)}|^{nq} \right)^{\frac{1}{q}} \leq W^{\frac{1}{q}} \|x\|_{\ell_p}^n.$$

Thus, $\|P\| \leq W^{\frac{1}{q}}$ and considering $\tilde{x} = W^{-\frac{s}{p}} (w_j^{\frac{s}{p}})_j \in S_{\ell_p}$, we obtain that $\|P\| = W^{\frac{1}{q}} = \|w\|_{\ell_s}^{\frac{1}{q}}$.

Finally, suppose that $n < \frac{p}{q}$ and $w \notin \ell_s$. Then, there exists $(b_j)_j \in \ell_{\frac{p}{nq}}$ with $b_1 \geq b_2 \geq b_3 \geq \dots \geq 0$ such that the series $\sum_{j=1}^{\infty} w_j b_j$ does not converge. Taking $\tilde{x} \in \ell_p$, $\tilde{x} = (b_j^{\frac{1}{nq}})_j$ we have that $P(\tilde{x}) = (b_j^{\frac{1}{q}})_j \notin d(w, q)$. Now, the proof is complete. \square

Proposition 3.8. $\mathcal{P}_w(^n\ell_p, d(w, q)) = \mathcal{P}(^n\ell_p, d(w, q))$ if and only if $n < \frac{p}{q}$ and $w \notin \ell_s$, with $s = (\frac{p}{nq})^*$.

Proof. By the previous proposition, whenever $n \geq \frac{p}{q}$ or $n < \frac{p}{q}$ and $w \in \ell_s$, $s = (\frac{p}{nq})^*$, the polynomial $P(x) = (x_j^n)_j$ belongs to $\mathcal{P}(^n\ell_p, d(w, q))$ and fails to be weakly continuous on bounded sets.

For the converse, by Remark 3.6 and Proposition 3.7, we have that it does not exist $P \in \mathcal{P}(^n\ell_p, d(w, q))$ such that $P(e_j) = f_j$, for every j . Finally, as $n < \frac{p}{q} \leq p \leq \text{cd}(\ell_p)$, all the hypothesis of Proposition 3.5 are fulfilled. Therefore, $\mathcal{P}(^n\ell_p, d(w, q)) = \mathcal{P}_{wsc0}(^n\ell_p, d(w, q))$. Now, by Proposition 3.4, $\mathcal{P}(^n\ell_p, d(w, q)) =$

$\mathcal{P}_{wsc}({}^n\ell_p, d(w, q))$ and the result follows from [8], since weakly sequentially continuous polynomials and weakly continuous polynomials on bounded sets coincide on ℓ_p . \square

Let $n = \text{cd}(\ell_p, d(w, q))$. Taking into account that for every $k < n$, any polynomial in $\mathcal{P}({}^k\ell_p, d(w, q))$ is weakly continuous on bounded sets, from the last proposition we derive that there are two possible values for n :

- (I) $\frac{p}{q} \leq n < \frac{p}{q} + 1$ and $w \notin \ell_{(\frac{p}{(n-1)q})^*}$, or
- (II) $n < \frac{p}{q}$ and $w \in \ell_{(\frac{p}{nq})^*} \setminus \ell_{(\frac{p}{(n-1)q})^*}$.

Theorem 3.9. *Let $n = \text{cd}(\ell_p, d(w, q))$.*

- (a) *If n and w satisfy condition (I) above, then*
 - $\mathcal{P}_w({}^n\ell_p, d(w, q))$ *is an M-ideal in $\mathcal{P}({}^n\ell_p, d(w, q))$, and*
 - $\mathcal{P}_w({}^k\ell_p, d(w, q))$ *is not a semi M-ideal in $\mathcal{P}({}^k\ell_p, d(w, q))$, for all $k > n$.*
- (b) *If n and w satisfy condition (II) above, then $\mathcal{P}_w({}^k\ell_p, d(w, q))$ is not a semi M-ideal in $\mathcal{P}({}^k\ell_p, d(w, q))$, for all $k \geq n$.*

Proof. Suppose n and w satisfy condition (I) above. Then, $n = \text{cd}(\ell_p, d(w, q)) \geq \frac{p}{q}$ and $\text{cd}(\ell_p)$ is the integer number satisfying $p \leq \text{cd}(\ell_p) < p + 1$. If $n < \text{cd}(\ell_p)$, the hypothesis of Theorem 2.5 are fulfilled. If $n = \text{cd}(\ell_p)$ we may apply Proposition 2.2. In both cases the conclusion follows.

Now, take $k > \text{cd}(\ell_p, d(w, q))$. According to Proposition 3.1, the result is proven if we find polynomials $P \in \mathcal{P}({}^k\ell_p, d(w, q))$ and $Q \in \mathcal{P}_w({}^k\ell_p, d(w, q))$ such that there exists $\delta > 0$ with $\|P\| = \|Q\|$ and $\|\pi^m P + Q\| \geq \delta > \|P\|$, for all m .

By Proposition 3.7, as $k - 1 \geq \frac{p}{q}$, the mapping $R(x) = (x_j^{k-1})_{j \geq 2}$ is a well defined norm one polynomial from ℓ_p to $d(w, q)$. Then, $P(x) = e_1^*(x)R(x)$ belongs to $\mathcal{P}({}^k\ell_p, d(w, q))$. In order to compute its norm, take x so that $\|x\|_{\ell_p} = 1$,

$$\|P(x)\|_{d(w,q)} = |x_1| \|R(x)\|_{d(w,q)} \leq |x_1| \|(x_j)_{j \geq 2}\|_{\ell_p}^{k-1} \leq \left(\frac{1}{k}\right)^{\frac{1}{p}} \left(1 - \frac{1}{k}\right)^{\frac{k-1}{p}},$$

where the last inequality was shown in the proof of Theorem 3.2. Now, with $\tilde{x} = \left(\frac{1}{k}\right)^{\frac{1}{p}} e_1 + \left(1 - \frac{1}{k}\right)^{\frac{1}{p}} e_2 \in S_{\ell_p}$ we have that $P(\tilde{x}) = \left(\frac{1}{k}\right)^{\frac{1}{p}} \left(1 - \frac{1}{k}\right)^{\frac{k-1}{p}} e_1$, whence $\|P\| = \left(\frac{1}{k}\right)^{\frac{1}{p}} \left(1 - \frac{1}{k}\right)^{\frac{k-1}{p}}$.

Let Q be the weakly continuous on bounded sets polynomial given by $Q = \|P\| (e_1^*)^k \cdot e_1$. Then, $\|Q\| = \|P\|$ and $\tilde{x} = \left(\frac{1}{k}\right)^{\frac{1}{p}} e_1 + \left(1 - \frac{1}{k}\right)^{\frac{1}{p}} e_{m+2}$, for $m \geq 1$, is a norm one vector so that

$$\|(\pi^m P + Q)(\tilde{x})\|_{d(w,q)} = \|P\| \left\| e_{m+1} + \left(\frac{1}{k}\right)^{\frac{k}{p}} e_1 \right\|_{d(w,q)} = \|P\| \left(1 + w_2 \left(\frac{1}{k}\right)^{\frac{kq}{p}}\right)^{\frac{1}{q}} > \|P\|,$$

which completes the proof of (a).

To prove (b), take n and w satisfying condition (II) and take $k \geq n$. Let us denote $s = \left(\frac{p}{nq}\right)^* = \frac{p}{p-nq}$ and $W = \|w\|_{\ell_s}$. By Proposition 3.7(b), the n -homogeneous

polynomial $R(x) = (x_j^n)_j$ satisfies

$$\|R(\tilde{x})\|_{d(w,q)} = \|R\| = W^{\frac{1}{q}}, \quad \text{where } \tilde{x} = W^{-\frac{s}{p}}(w_j^{\frac{s}{p}})_j.$$

Observe that $x^* = (w_j^{\frac{s}{p^*}})_j$ belongs to ℓ_{p^*} and, as a continuous functional, it also attains its norm at \tilde{x} :

$$x^*(\tilde{x}) = \|x^*\| = W^{\frac{s}{p^*}}.$$

Now we are ready to construct two polynomials P and Q fulfilling the statement of Proposition 3.1. Let $P \in \mathcal{P}({}^k\ell_p, d(w, q))$ and $Q \in \mathcal{P}_w({}^k\ell_p, d(w, q))$ be given by

$$P(x) = x^*(x)^{k-n}R(x) \quad \text{and} \quad Q(x) = W^{\frac{1}{q} - \frac{sn}{p^*}}x^*(x)^k e_1.$$

It is easy to see that

$$\|P(\tilde{x})\|_{d(w,q)} = \|P\| = W^r = \|Q(\tilde{x})\|_{d(w,q)} = \|Q\|, \quad \text{where } r = \frac{s(k-n)}{p^*} + \frac{1}{q}.$$

Finally,

$$\begin{aligned} \|\pi^m P + Q\| &\geq \|(\pi^m P + Q)(\tilde{x})\|_{d(w,q)} = \left\| W^r e_1 + W^{\frac{s(k-n)}{p^*}} \sum_{j=m+1}^{\infty} \tilde{x}_j^n e_j \right\|_{d(w,q)} \\ &= W^r \left[1 + W^{-1} \sum_{j=2}^{\infty} w_j |\tilde{x}_{m-1+j}|^{nq} \right]^{\frac{1}{q}} > W^r = \|P\|. \end{aligned}$$

This completes the proof of the theorem. □

4. Polynomial property (M)

Property (M) was introduced by Kalton in [26]. It is a geometric property relating the norm of the traslation by a weakly null net of any two elements of the space. Namely, a Banach space X has property (M) if for any $x, \tilde{x} \in X$ such that $\|x\| \leq \|\tilde{x}\|$, and any bounded weakly null net $(x_\alpha)_\alpha$ in X , it holds that $\limsup \|x + x_\alpha\| \leq \limsup \|\tilde{x} + x_\alpha\|$. An operator version of this property was given in [27]. Later on, in [16], it is extended to the scalar-valued polynomial context. In all these cases, these properties have incidence in the correspondent M -ideal problems. To study M -structures in spaces of vector-valued polynomials, we consider a suitable property (M), which is the result of a natural combination of the definitions given for operators and scalar-valued polynomials. Before going on, let us state the vector-valued versions of [16, Lemma 3.1 and Theorem 3.2].

Lemma 4.1. *If $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$ then, for each $P \in \mathcal{P}({}^n E, F)$ there exists a bounded net $\{P_\alpha\}_\alpha \subset \mathcal{P}_w({}^n E, F)$ such that $P_\alpha(x) \rightarrow P(x)$, for all $x \in E$.*

Proof. Fix $P \in \mathcal{P}({}^nE, F)$. By [24, Remark I.1.13], we may consider $\{Q_\alpha\}_\alpha$ a bounded net in $\mathcal{P}_w({}^nE, F)$ such that $Q_\alpha \rightarrow P$ in the topology $\sigma(\mathcal{P}({}^nE, F), \mathcal{P}_w({}^nE, F)^*)$.

Since $e_x \otimes y^*$ belongs to $\mathcal{P}_w({}^nE, F)^*$, $y^*(Q_\alpha(x)) = \langle e_x \otimes y^*, Q_\alpha \rangle \rightarrow \langle e_x \otimes y^*, P \rangle = y^*(P(x))$, for all $x \in E$ and all $y^* \in F^*$. This says that $Q_\alpha(x) \xrightarrow{w} P(x)$, for all $x \in E$, which can be described, in analogy to the operator setting, as $Q_\alpha \rightarrow P$ in the *WPT*, the “weak polynomial topology”.

We can also consider on $\mathcal{P}({}^nE, F)$ the “strong polynomial topology”, *SPT*, naturally meaning pointwise convergence of nets. Both topologies, the *WPT* and the *SPT*, are locally convex and have the same continuous functionals (the proof of [19, Theorem VI.1.4] works also for polynomials). Thus, as in the linear case, we derive that the closure of any convex set in the strong polynomial topology coincides with its closure in the weak polynomial topology.

Then, we may find P_α , a convex combination of Q_α , converging pointwise to P . \square

As a consequence of [32, Proposition 2.3] and the previous lemma, we have the following result which can be proved analogously to [32, Theorem 3.1]:

Theorem 4.2. *Let E and F be Banach spaces. The following are equivalent:*

- (i) $\mathcal{P}_w({}^nE, F)$ is an M -ideal in $\mathcal{P}({}^nE, F)$.
- (ii) For all $P \in \mathcal{P}({}^nE, F)$ there exists a net $\{P_\alpha\}_\alpha \subset \mathcal{P}_w({}^nE, F)$ such that $P_\alpha(x) \rightarrow P(x)$, for all $x \in E$ and

$$\limsup \|Q + P - P_\alpha\| \leq \max\{\|Q\|, \|Q\|_{es} + \|P\|\}, \text{ for all } Q \in \mathcal{P}({}^nE, F).$$

- (iii) For all $P \in \mathcal{P}({}^nE, F)$ there exists a net $\{P_\alpha\}_\alpha \subset \mathcal{P}_w({}^nE, F)$ such that $P_\alpha(x) \rightarrow P(x)$, for all $x \in E$ and

$$\limsup \|Q + P - P_\alpha\| \leq \max\{\|Q\|, \|P\|\}, \text{ for all } Q \in \mathcal{P}_w({}^nE, F).$$

Now we state the property (M) for a vector-valued polynomial.

Definition 4.3. Let $P \in \mathcal{P}({}^nE, F)$ with $\|P\| \leq 1$. We say that P has **property (M)** if for all $u \in E$, $v \in F$ with $\|v\| \leq \|u\|^n$ and for every bounded weakly null net $\{x_\alpha\}_\alpha \subset E$, it holds that

$$\limsup_\alpha \|v + P(x_\alpha)\| \leq \limsup_\alpha \|u + x_\alpha\|^n.$$

Note that every $P \in \mathcal{P}_w({}^nE, F)$ with $\|P\| \leq 1$ has property (M) . Analogously to [27, Lemma 6.2], we can prove:

Lemma 4.4. *Let $P \in \mathcal{P}({}^nE, F)$ with $\|P\| \leq 1$. If P has property (M) then for all nets $\{u_\alpha\}_\alpha$ and $\{v_\alpha\}_\alpha$ contained in compact sets of E and F respectively, with $\|v_\alpha\| \leq \|u_\alpha\|^n$ and for every bounded weakly null net $\{x_\alpha\}_\alpha \subset E$, it holds that*

$$\limsup_\alpha \|v_\alpha + P(x_\alpha)\| \leq \limsup_\alpha \|u_\alpha + x_\alpha\|^n.$$

Definition 4.5. We say that a pair of Banach spaces (E, F) has the n -**polynomial property** (M) if every $P \in \mathcal{P}({}^n E, F)$ with $\|P\| \leq 1$ has property (M) .

The next two results can be proved mimicking the proofs of Proposition 3.7 and Theorem 3.9 of [16].

Proposition 4.6. *If $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$ and $n = \text{cd}(E, F)$ then (E, F) has the n -polynomial property (M) .*

Theorem 4.7. *Let E and F be Banach spaces and suppose that there exists a net of compact operators $\{K_\alpha\}_\alpha \in \mathcal{K}(E)$ satisfying the following two conditions:*

- $K_\alpha(x) \rightarrow x$, for all $x \in E$ and $K_\alpha^*(x^*) \rightarrow x^*$, for all $x^* \in E^*$.
- $\|Id - 2K_\alpha\| \xrightarrow{\alpha} 1$.

Suppose also that $n = \text{cd}(E, F)$. Then, $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$ if and only if (E, F) has the n -polynomial property (M) .

Sometimes it is possible to infer M -structures in the space of linear continuous operators from the existence of geometric structures on the underlying space. For instance, it is proved in [24, Theorem VI.4.17] that $\mathcal{K}(E)$ is an M -ideal in $\mathcal{L}(E)$ if and only if E has property (M) and satisfies both conditions of the theorem above. A similar result [16, Theorem 3.9] is obtained in the scalar-valued polynomial setting for $n = \text{cd}(E)$ using the polynomial property (M) . The following proposition (which is the vector-valued polynomial version of [24, Lemma VI.4.14] and [16, Proposition 3.10]) paves the way to connect the linear M -structure with M -ideals in vector valued polynomial spaces.

Proposition 4.8. *Let E and F be Banach spaces and $n = \text{cd}(E, F) < \text{cd}(E)$. If E and F have the property (M) , then (E, F) has the n -polynomial property (M) .*

Proof. Let $P \in \mathcal{P}({}^n E, F)$ with $\|P\| = 1$. Fix $u \in E$, $v \in F$ with $\|v\| \leq \|u\|^n$ and a bounded weakly null net $\{x_\alpha\}_\alpha \subset E$. We want to prove that

$$\limsup_{\alpha} \|v + P(x_\alpha)\| \leq \limsup_{\alpha} \|u + x_\alpha\|^n.$$

Given $\varepsilon > 0$, take $x \in S_E$ such that $\|P(x)\| > 1 - \varepsilon$ and $\tilde{x} = \|v\|^{\frac{1}{n}} x$. Then, $(1 - \varepsilon)\|v\| < \|P(\tilde{x})\| \leq \|\tilde{x}\| \leq \|u\|$. As $n < \text{cd}(E)$, every scalar valued polynomial in $\mathcal{P}({}^n E)$ is weakly continuous on bounded sets. Then, P is weak-to-weak continuous and $P(x_\alpha) \xrightarrow{w} 0$. Therefore, since F has property (M) ,

$$\begin{aligned} \limsup_{\alpha} \|(1 - \varepsilon)v + P(x_\alpha)\| &\leq \limsup_{\alpha} \|P(\tilde{x}) + P(x_\alpha)\| \\ &= \limsup_{\alpha} \|P(\tilde{x} + x_\alpha)\| \\ &\leq \limsup_{\alpha} \|\tilde{x} + x_\alpha\|^n \\ &\leq \limsup_{\alpha} \|u + x_\alpha\|^n, \end{aligned}$$

where the last inequality holds since E has property (M) . Now, letting $\varepsilon \rightarrow 0$ we obtain the desired inequality.

If $\|P\| < 1$, the result follows from the previous case through the following convex combination

$$v + P(x_\alpha) = \frac{1 + \|P\|}{2} \left(v + \frac{P}{\|P\|}(x_\alpha) \right) + \frac{1 - \|P\|}{2} \left(v - \frac{P}{\|P\|}(x_\alpha) \right). \quad \square$$

Now we can lift M -structures from the linear to the vector-valued polynomial context. This is done for the particular case of n -homogeneous polynomials when n is the critical degree of the pair (E, F) and it is strictly less than the critical degree of the domain space E . We do not know if the result remains true even for the case $n = \text{cd}(E, F) = \text{cd}(E)$.

Corollary 4.9. *Let E and F be Banach spaces and $n = \text{cd}(E, F) < \text{cd}(E)$. If $\mathcal{K}(E)$ is an M -ideal in $\mathcal{L}(E)$ and F has property (M) , then $\mathcal{P}_w({}^n E, F)$ is an M -ideal in $\mathcal{P}({}^n E, F)$.*

Proof. If $\mathcal{K}(E)$ is an M -ideal in $\mathcal{L}(E)$, appealing to [24, Theorem VI.4.17], E has property (M) and we may find $\{K_\alpha\}_\alpha \subset \mathcal{K}(E)$ a net of compact operators satisfying both conditions of Theorem 4.7. By Proposition 4.8, (E, F) has the n -polynomial property (M) . Now, we may apply Theorem 4.7 to derive the result. \square

We finish this section applying the previous result to give some examples of M -ideals of polynomials between Bergman and ℓ_p spaces.

Example 4.10. The Bergman space B_p is the space of all holomorphic functions in $L_p(\mathbb{D}, dxdy)$, where \mathbb{D} is the complex disc. If $1 < p < \infty$, B_p is isomorphic to ℓ_p [33, Theorem III.A.11] and so, for $1 < p, q < \infty$,

$$\text{cd}(\ell_p, \ell_q) = \text{cd}(\ell_p, B_q) = \text{cd}(B_p, \ell_q) = \text{cd}(B_p, B_q).$$

Since, by [27, Corollary 4.8], $\mathcal{K}(B_p)$ is an M -ideal in $\mathcal{L}(B_p)$, we obtain from Corollary 4.9, that, if $n = \text{cd}(\ell_p, \ell_q) < \text{cd}(\ell_p)$, then:

- $\mathcal{P}_w({}^n \ell_p, B_q)$ is an M -ideal in $\mathcal{P}({}^n \ell_p, B_q)$.
- $\mathcal{P}_w({}^n B_p, \ell_q)$ is an M -ideal in $\mathcal{P}({}^n B_p, \ell_q)$.
- $\mathcal{P}_w({}^n B_p, B_q)$ is an M -ideal in $\mathcal{P}({}^n B_p, B_q)$.

Acknowledgements. We would like to thank the anonymous referee for his/her comments that improve the presentation of the article.

References

[1] M. D. Acosta: Denseness of norm attaining mappings, RACSAM, Rev. R. Acad. Cienc. Exactas Fís. Nat., Ser. A Mat. 100(1-2) (2006) 9-30.
 [2] M. D. Acosta, F. J. Aguirre, R. Payá: There is no bilinear Bishop-Phelps theorem, Isr. J. Math. 93 (1996) 221-227.

- [3] E. Alfsen, E. Effros: Structure in real Banach spaces. I, II. *Ann. of Math.* (2)96 (1972) 98–128; *Ann. of Math.* (2)96 (1972) 129–173.
- [4] R. Alencar, K. Floret: Weak-strong continuity of multilinear mappings and the Pelczynski-Pitt theorem, *J. Math. Anal. Appl.* 206(2) (1997) 532–546.
- [5] R. Aron, P. Berner: A Hahn-Banach extension theorem for analytic mappings, *Bull. Soc. Math. Fr.* 106 (1978) 3–24.
- [6] R. Aron, V. Dimant: Sets of weak sequential continuity for polynomials, *Indag. Math., New Ser.* 13(3) (2002) 287–299.
- [7] R. Aron, D. García, M. Maestre: On norm attaining polynomials, *Publ. Res. Inst. Math. Sci.* 39(1) (2003) 165–172.
- [8] R. Aron, C. Hervés, M. Valdivia: Weakly continuous mappings on Banach spaces, *J. Funct. Anal.* 52 (1983) 189–204.
- [9] R. Aron, J. B. Prolla: Polynomial approximation of differentiable functions on Banach spaces, *J. Reine Angew. Math.* 313 (1980) 195–216.
- [10] R. M. Aron, M. Schottenloher: Compact holomorphic mappings on Banach spaces and the approximation property, *J. Funct. Anal.* 21(1) (1976) 7–30.
- [11] E. Asplund: Farthest points in reflexive locally uniformly rotund Banach spaces, *Isr. J. Math.* 4 (1966) 213–216.
- [12] P. Bandyopadhyay, B.-L. Lin, T. S. S. R. K. Rao: Ball remotal subspaces of Banach spaces, *Colloq. Math.* 114(1) (2009) 119–133.
- [13] C. Boyd, R. A. Ryan: Bounded weak continuity of homogeneous polynomials at the origin, *Arch. Math. (Basel)* 71(3) (1998) 211–218.
- [14] D. Carando, S. Lassalle: E' and its relation with vector-valued functions on E , *Ark. Mat.* 42(2) (2004) 283–300.
- [15] Y. S. Choi, S. G. Kim: Norm or numerical radius attaining multilinear mappings and polynomials, *J. Lond. Math. Soc., II. Ser.* 54(1) (1996) 135–147.
- [16] V. Dimant: M -ideals of homogeneous polynomials, *Stud. Math.* 202 (2011) 81–104.
- [17] V. Dimant, R. Gonzalo: Block diagonal polynomials, *Trans. Amer. Math. Soc.* 353(2) (2001) 733–747.
- [18] S. Dineen: *Complex Analysis on Infinite-Dimensional Spaces*, Springer Monographs in Mathematics, Springer, London (1999).
- [19] N. Dunford, J. T. Schwartz: *Linear Operators. Part I: General Theory*, Wiley, New York (1958).
- [20] M. Edelstein: Farthest points of sets in uniformly convex Banach spaces, *Isr. J. Math.* 4 (1966) 171–176.
- [21] M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant, V. Zizler: *Functional Analysis and Infinite-Dimensional Geometry*, CMS Books in Mathematics / Ouvrages de Mathématiques de la SMC 8, Springer, New York (2001).
- [22] M. González, J. M. Gutiérrez: The polynomial property (V), *Arch. Math.* 75(4) (2000) 299–306.
- [23] R. Gonzalo, J. A. Jaramillo: Compact polynomials between Banach spaces, *Proc. R. Ir. Acad., Sect. A* 95(2) (1995) 213–226.

- [24] P. Harmand, D. Werner, W. Werner: *M-Ideals in Banach Spaces and Banach Algebras*, Lecture Notes in Mathematics 1547, Springer, Berlin (1993).
- [25] J. Hennefeld: *M-ideals, HB-subspaces, and compact operators*, Indiana Univ. Math. J. 28(6) (1979) 927–934.
- [26] N. J. Kalton: *M-ideals of compact operators*, Ill. J. Math. 37(1) (1993) 147–169.
- [27] N. J. Kalton, D. Werner: *Property (M), M-ideals, and almost isometric structure of Banach spaces*, J. Reine Angew. Math. 461 (1995) 137–178.
- [28] A. Lima: *M-ideals of compact operators in classical Banach spaces*, Math. Scand. 44(1) (1979) 207–217.
- [29] J. Mujica: *Complex Analysis in Banach Spaces*, North-Holland Mathematics Studies 120, North-Holland, Amsterdam (1986).
- [30] E. Oja: *On M-ideals of compact operators and Lorentz sequence spaces*, Proc. Est. Acad. Sci., Phys. Math. 40(1) (1991) 31–36.
- [31] E. Oja, D. Werner: *Remarks on M-ideals of compact operators on $X \oplus_p X$* , Math. Nachr. 152 (1991) 101–111.
- [32] D. Werner: *M-ideals and the “basic inequality”*, J. Approximation Theory 76(1) (1994) 21–30.
- [33] P. Wojtaszczyk: *Banach Spaces for Analysts*, Cambridge Studies in Advanced Mathematics 25, Cambridge University Press, Cambridge (1991).