

An example of a solvable Lie algebra

Michael Wüstner

We shall consider a solvable real Lie algebra \mathfrak{g} which is linearly spanned by the set $\text{comp } \mathfrak{g}$ of its compact elements and use the notation of [1]. In particular, the diagram in [1] below Proposition 2.10 gives a good orientation on the subalgebras involved in our discussion. It was shown in [1] that a Cartan algebra in a real Lie algebra spanned by its compact elements has nilpotent class ≤ 2 . It was left open whether there are algebras in which the Cartan algebras are nonabelian. In the following we present an example of dimension 13 showing that this can happen.

Let \mathfrak{g} be a real vector space with a basis

$$\{u, x, y, z, t, e_1, e_2, f_1, f_2, g_1, g_2, h_1, h_2\}$$

and define the following relations on the basis elements:

$$\begin{aligned} [u, a_1] &= a_2, & [u, a_2] &= -a_1 & \text{for } a &= e, f, g, h, \\ [e_1, e_2] &= x, & [f_1, f_2] &= y, \\ [e_1, f_2] &= -t, & [e_2, f_1] &= t, \\ [x, f_i] &= h_i, & [y, e_i] &= g_i & \text{for } i &= 1, 2, & [x, y] &= z, \\ [t, e_i] &= -h_i, & [t, f_i] &= -g_i & \text{for } i &= 1, 2, \\ [e_1, g_2] &= -\frac{1}{2}z, & [e_2, g_1] &= \frac{1}{2}z, & [f_1, h_2] &= \frac{1}{2}z, & [f_2, h_1] &= -\frac{1}{2}z. \end{aligned}$$

All other brackets between basis elements are set equal to 0. Bilinear extension yield a bilinear product $[\cdot, \cdot]: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$.

We define on \mathfrak{g} the structure of a graded vector space by setting $\mathfrak{g}^0 = \mathbb{R} \cdot u$, $\mathfrak{g}^1 = \text{span}\{e_1, e_2, f_1, f_2\}$, $\mathfrak{g}^2 = \text{span}\{x, y, t\}$, $\mathfrak{g}^3 = \text{span}\{g_1, g_2, h_1, h_2\}$, $\mathfrak{g}^4 = \mathbb{R} \cdot z$. It is readily verified that $[\mathfrak{g}^m, \mathfrak{g}^n] \subseteq \mathfrak{g}^{m+n}$. Thus \mathfrak{g} is a graded algebra with respect to the bracket.

Lemma. \mathfrak{g} is a solvable Lie algebra.

Proof. We verify the validity of the Jacobi identity by straightforward, but tedious calculation. I have written a program which executes the necessary verifications. The gradation permits us a quick inspection of the solvability of \mathfrak{g} . ■

We verify quickly that the subspace $\mathfrak{h} = \text{span}\{u, x, y, z, t\}$ is a Cartan subalgebra and $\mathfrak{t} \stackrel{\text{def}}{=} \mathfrak{h} \cap \text{comp } \mathfrak{g} = \text{span}\{u, z\}$. Also, the sum of the weight spaces with respect to \mathfrak{t} is $\mathfrak{t}^+ = \text{span}\{e_1, e_2, f_1, f_2, g_1, g_2, h_1, h_2\}$ (cf. [1]), and $\mathfrak{g}' = \mathcal{C}^\infty = \langle \mathfrak{t}^+ \rangle = \text{span}\{e_1, e_2, f_1, f_2, g_1, g_2, h_1, h_2, t, x, y, z\}$. It follows that $\mathfrak{e} \stackrel{\text{def}}{=} \mathfrak{h} \cap \mathfrak{g}' = \text{span}\{x, y, t, z\}$. Further, in the notation of [1], we have $\mathfrak{z} = \mathbb{R} \cdot z = \mathfrak{j}$ and may take $\mathfrak{e}_1 = \text{span}\{x, y, t\}$ as a complement of \mathfrak{j} in \mathfrak{e} , and $\mathfrak{t}_1 = \mathbb{R} \cdot u$ as a complement

of \mathfrak{z} in \mathfrak{t} . With these choices we obtain $\mathfrak{g}' = \mathfrak{e}_1 \oplus \mathfrak{t}^+$. Therefore, we have $\dim \mathfrak{t} = 2$, $\dim \mathfrak{h} = 5$, $\dim \mathfrak{e} = 4$, $\dim \mathfrak{e}_1 = 3$, $\dim \mathfrak{t}^+ = 8$, $\dim \mathfrak{g}' = \dim \langle \mathfrak{t}^+ \rangle = 12$.

The following diagram records the gradation and the relevant subalgebras:

| degree | \mathfrak{h} | | \mathfrak{t}^+ | |
|--------|----------------|------------------|------------------|-------------------|
| | \mathfrak{t} | \mathfrak{e}_1 | | |
| 0 | u | | | |
| 1 | | | e_1 | e_2 f_1 f_2 |
| 2 | | x y t | | |
| 3 | | | g_1 g_2 | h_1 h_2 |
| 4 | z | | | |

The following equations will show that $\mathfrak{g} = \text{span}(\text{comp}(\mathfrak{g}))$. From [1], Lemma 2.2(iii) we know that $\text{comp} = \Gamma \cdot \mathfrak{t}$. We will be finished if we can show that \mathfrak{g} is spanned by elements of the form $e^{ad w}(u)$ with $w \in \mathcal{C}^\infty$. Now we compute:

$$\begin{aligned}
 e^{ad(e_1+e_2)}(u) &= u - e_2 + e_1 - 2x, \\
 e^{ad(e_1-e_2)}(u) &= u - e_2 - e_1 - 2x, \\
 e^{ad(e_2-e_1)}(u) &= u + e_1 + e_2 - 2x, \\
 e^{ad(f_1+f_2)}(u) &= u - f_2 + f_1 - 2y, \\
 e^{ad(f_1-f_2)}(u) &= u - f_2 - f_1 - 2y, \\
 e^{ad(f_2-f_1)}(u) &= u + f_1 + f_2 - 2y, \\
 e^{ad(e_1+f_1)}(u) &= u - e_2 - f_2 - x - y + 3g_1 + 3h_1 + 2t, \\
 e^{ad(a_1)}(u) &= u - a_2, \\
 e^{ad(a_2)}(u) &= u + a_1
 \end{aligned}$$

for $a = g, h$.

We see that the span of these elements is \mathfrak{g} and this shows that $\mathfrak{g} = \text{span comp}$.

References

- [1] Hofmann, K.,H., *Compact elements in solvable real Lie algebras*, Seminar Sophus Lie **2** (1992), this volume.

Fachbereich Mathematik
Technische Hochschule
Schloßgartenstr. 7
D-6100 Darmstadt

Received May 12, 1992