

# A Variational Principle in Reflexive Spaces with Kadec-Klee Norm

**Marián Fabian\***

*Mathematical Institute, Czech Academy of Sciences,  
Žitná 25, 115 67 Praha 1, Bohemia  
fabian@math.cas.cz*

**Julian Revalski†**

*on leave from: Institute of Mathematics and Informatics,  
Bulgarian Academy of Sciences, Acad. G. Bonchev Street, Block 8, 1113 Sofia, Bulgaria,  
revalski@math.bas.bg*

*Laboratoire AOC, Département de Mathématiques et Informatique,  
Université des Antilles et de la Guyane, BP 592, Campus de Fouillole,  
97157 Pointe-à-Pitre, Guadeloupe, France  
julian.revalski@univ-ag.fr*

Received: May 21, 2007

Revised manuscript received: February 6, 2008

We prove a variational principle in reflexive Banach spaces  $X$  with Kadec-Klee norm, which asserts that any Lipschitz (or any proper lower semicontinuous bounded from below extended real-valued) function in  $X$  can be perturbed with a parabola in such a way that the perturbed function attains its infimum (even more can be said – the infimum is well-posed). In addition, we have genericity of the points determining the parabolas. We prove also that the validity of such a principle actually characterizes the reflexive spaces with Kadec-Klee norm. This principle turns out to be an analytic counterpart of a result of K.-S. Lau on nearest points.

*2000 Mathematics Subject Classification:* 46B20, 49J52

## 1. Introduction and main result

In 1961, Bishop and Phelps proved a lemma on supporting closed sets by cones [1]. An analytic counterpart of it, on supporting functions, is the Ekeland variational principle [6], stating that any (extended real-valued) lower semicontinuous bounded from below function in a complete metric space can be perturbed by a distance-like function in such a way that the perturbed function attains its minimum. In 1974, Phelps [14] proved that in a Banach space which has a geometric property, called Radon-Nikodým property, every nonempty bounded closed convex set is the closed convex hull of its strongly exposed points. An analytic result related to this geometric fact is the Stegall's variational principle [17, 18] which shows that in such a space one can perturb a lower

\*The first author was supported by the Institutional Research Plan of the Academy of Sciences of Czech Republic AVOZ 101 905 03 and by GAČR 201/07/0394 and also by Université des Antilles et de la Guayane, where he was a visiting professor for one month.

†The second author was supported by Bulgarian National Fund for Scientific Research under contract DO 02-360/2008.

semicontinuous bounded from below function (with domain in a closed bounded convex set) by a linear continuous functional, in order to obtain a perturbation which attains its minimum.

The aim of this note is to go on in this vein and to show one more example of such a correspondence between a geometrical and an analytical statement. In 1978, Lau [13] proved that in a reflexive Banach space  $(X, \|\cdot\|)$ , with Kadec-Klee norm, for every nonempty closed set  $C \subset X$  there are residually many points in  $X \setminus C$  possessing a nearest point in  $C$ . An analytic counterpart of this result is the following theorem. And, likewise, as in the previous cases, we can formulate it in the form of equivalence(s).

**Theorem 1.1.** *For a Banach space  $(X, \|\cdot\|)$  the following statements are equivalent:*

- (i) *The space  $X$  is reflexive and its norm  $\|\cdot\|$  is Kadec-Klee.*
- (ii) *For every Lipschitzian function  $f : X \rightarrow (-\infty, +\infty)$  there exists a residual set  $\Omega \subset X$  such that for every  $u \in \Omega$  there is a residual set  $R_u \subset (0, +\infty)$  such that for each  $c \in R_u$  the problem of minimizing  $f + c\|\cdot - u\|^2$  is well-posed.*
- (ii') *For every proper lower semicontinuous function  $f : X \rightarrow (-\infty, +\infty]$ , with  $\inf f > -\infty$ , there exists a residual set  $\Omega \subset X$  such that for every  $u \in \Omega$  there are  $a_u \in (0, +\infty]$  and a residual set  $R_u \subset (0, a_u)$  such that for each  $c \in R_u$  the problem of minimizing the function  $f + c\|\cdot - u\|^2$  is well-posed.*
- (iii) *For every  $f$  as in (ii) there exists a dense set  $D \subset X$  such that for every  $u \in D$  the function  $f + \|\cdot - u\|^2$  attains its infimum.*
- (iii') *For every  $f$  as in (ii') there exists a dense set  $D \subset X$  such that for every  $u \in D$  there is  $c > 0$  so that the function  $f + c\|\cdot - u\|^2$  attains its infimum.*

We follow the usual notation in Banach space theory, see, e.g. [10]. Let us recall that the norm  $\|\cdot\|$  on  $X$  is called *Kadec-Klee* if  $\lim_{n \rightarrow \infty} \|x_n - x_0\| = 0$  whenever a sequence  $x_0, x_1, x_2, \dots \in X$  satisfies  $\|x_n\| \rightarrow \|x_0\|$  and  $x_n \rightarrow x_0$  weakly. In Russian literature, a reflexive Banach space, with Kadec-Klee norm, is called a *Efimov-Stečkin space*. A set in a Baire space, in particular, in a Banach space, is called *residual* if its complement is the union of countably many nowhere dense sets. An extended real-valued function  $g : X \rightarrow (-\infty, +\infty]$  is called *proper* if its effective domain  $\text{dom } g = \{x \in X; g(x) < +\infty\}$  is nonempty. Given such a function  $g$ , with  $\inf g > -\infty$ , we say that *the problem of minimizing  $g$  is well-posed*, if every minimizing sequence  $(x_n)_{n \in \mathbb{N}}$  in  $X$  for  $g$  (i.e.  $g(x_n) \downarrow \inf g$ ) has a norm-cluster point  $x \in X$ ; hence  $g(x) = \inf g$  whenever  $g$  is at least lower semicontinuous. Obviously, if the problem to minimize  $g$  is well-posed, then the set of minimizers for  $g$  is compact.

In the proof, we shall need the following topological fact going back to Ulam and Kuratowski.

**Lemma 1.2.** *Let  $X$  be a Banach space and let  $O \subset X \times \mathbb{R}$  be an open subset such that its canonical projection, along  $\mathbb{R}$ , is all of  $X$ . Let  $E \subset O$  be a residual subset in  $O$ . Then there exists a residual subset  $\Omega \subset X$  such that for every  $x \in \Omega$  the set  $\{t \in \mathbb{R}; (x, t) \in E\}$  is residual in  $\{t \in \mathbb{R}; (x, t) \in O\}$ .*

**Proof.** Put  $F = E \cup [(X \times \mathbb{R}) \setminus O]$ . It is easy to check that  $F$  is residual in  $X \times \mathbb{R}$ . Indeed,

$$(X \times \mathbb{R}) \setminus F = [(X \times \mathbb{R}) \setminus E] \cap O = O \setminus E,$$

and the latter set is of the first Baire category in  $O$ , and hence in  $X \times \mathbb{R}$ . Now, by applying [12, §22.V, Corollary 1a], we find a residual set  $\Omega \subset X$  such that for every  $x \in \Omega$  the set  $\{t \in \mathbb{R}; (x, t) \in F\}$  is residual in  $\mathbb{R}$ . Hence, the set  $\{t \in \mathbb{R}; (x, t) \in E\}$  is residual in  $\{t \in \mathbb{R}; (x, t) \in O\}$ .  $\square$

**Proof of Theorem 1.1.** (i)  $\Rightarrow$  (ii). We will split the proof of this implication into several steps.

*Step 1.* Let  $f$  be as in (ii). Consider the (reflexive) space  $X \times \mathbb{R}$  and the (closed) set  $\text{epi } f = \{(x, t) \in X \times \mathbb{R}; f(x) \leq t\}$ . Let  $L > 0$  be a Lipschitzian constant of  $f$  and put

$$B = \{(x, t) \in X \times \mathbb{R}; |t| \leq -\|x\|^2 + L^2\}. \tag{1}$$

It is elementary to check that the set  $B$  is closed, convex, symmetric, bounded, and that it contains the origin  $(0, 0)$  in its interior. Let  $\|\cdot\|$  denote its Minkowski functional; this will be an equivalent norm on  $X \times \mathbb{R}$ .

For sure, this norm will have also the Kadec-Klee property. Indeed, for  $n = 0, 1, 2, \dots$  consider  $(x_n, t_n) \in X \times \mathbb{R}$  such that  $\|(x_n, t_n)\| = 1$ , and  $(x_n, t_n) \rightarrow (x_0, t_0)$  weakly as  $n \rightarrow \infty$ . Then, of course,  $x_n \rightarrow x_0$  weakly and  $t_n \rightarrow t_0$  as  $n \rightarrow \infty$ . Now, for all  $n = 0, 1, \dots$ , from the very definition of  $\|\cdot\|$ , we must have  $|t_n| + \|x_n\|^2 = L^2$  and hence

$$\|x_0\| \leq \liminf_{n \rightarrow \infty} \|x_n\| \leq \limsup_{n \rightarrow \infty} \|x_n\| = \lim_{n \rightarrow \infty} \sqrt{L^2 - |t_n|} = \sqrt{L^2 - |t_0|} = \|x_0\|.$$

Thus, the Kadec-Klee property of  $\|\cdot\|$  yields that  $\|x_n - x_0\| \rightarrow 0$  and so  $\|(x_n, t_n) - (x_0, t_0)\| \rightarrow 0$  as  $n \rightarrow \infty$ .

*Step 2.* Put  $O = (X \times \mathbb{R}) \setminus \text{epi } f$  and consider the distance function for  $\text{epi } f$

$$\text{dist}((x, t), \text{epi } f) := \inf \{\|(x, t) - (u, s)\|; (u, s) \in \text{epi } f\}, \quad (x, t) \in O.$$

We can immediately see that  $O$  is an open set and that its projection, along  $\mathbb{R}$ , is equal to all of  $X$ . By the mentioned result of Lau, [13, Theorem 4] (see also [2, Corollary 5.8]), we find a residual subset  $E$  of  $O$ , such that for every  $(x_0, t_0) \in E$ , if  $(y_1, s_1), (y_2, s_2), \dots \in \text{epi } f$ , and

$$\|(y_n, s_n) - (x_0, t_0)\| \longrightarrow \text{dist}((x_0, t_0), \text{epi } f) \quad \text{as } n \rightarrow \infty,$$

then the sequence  $((y_n, s_n))_{n \in \mathbb{N}}$  has a norm-cluster point (such a cluster point is evidently a nearest point to  $(x_0, t_0)$  in  $\text{epi } f$ ). In other words, the best approximation problem of any  $(x_0, t_0) \in E$  to  $\text{epi } f$  is well-posed. By the above Lemma applied for our  $E$  and  $O$ , we find the corresponding residual set  $\Omega \subset X$ . Fix, for a longer while, any  $x_0 \in \Omega$  and then find a residual set  $S \subset (-\infty, f(x_0))$  such that  $\{x_0\} \times S \subset E$ . Further, fix, for a longer while, any  $t_0 \in S$ . Let  $(x_1, t_1) \in \text{epi } f$  be a nearest point to  $(x_0, t_0)$  in  $\text{epi } f$ , i.e.,

$$(0 <) \rho := \|(x_1, t_1) - (x_0, t_0)\| \leq \|(x, t) - (x_0, t_0)\| \quad \text{for all } (x, t) \in \text{epi } f. \tag{2}$$

From the very definition of the norm  $\|\cdot\|$ , we have that

$$\rho|t_1 - t_0| + \|x_1 - x_0\|^2 = \rho^2 L^2. \tag{3}$$

Consider any  $\varepsilon \in (0, 1)$ . Then

$$\begin{aligned} & \|((x_1 - \varepsilon(x_1 - x_0), 2t_0 - t_1 - \varepsilon(t_0 - t_1)) - (x_0, t_0))\| \\ &= (1 - \varepsilon)\|(x_1 - x_0, t_0 - t_1)\| = (1 - \varepsilon)\rho < \rho, \end{aligned}$$

and so  $(x_1 - \varepsilon(x_1 - x_0), 2t_0 - t_1 - \varepsilon(t_0 - t_1)) \notin \text{epi } f$ , that is,

$$2t_0 - t_1 - \varepsilon(t_0 - t_1) < f(x_1 - \varepsilon(x_1 - x_0)).$$

Letting  $\varepsilon \downarrow 0$  in the last inequality we obtain  $2t_0 - t_1 \leq f(x_1)$ . However,  $(x_1, t_1) \in \text{epi } f$ , and so  $f(x_1) \leq t_1$ . Therefore  $2t_0 - t_1 \leq t_1$ , that is,  $t_0 \leq t_1$ . Thus (3) has the form

$$\rho(t_1 - t_0) + \|x_1 - x_0\|^2 = \rho^2 L^2. \quad (4)$$

We actually have that  $t_1 = f(x_1)$ . Indeed, assume that  $t_1 = f(x_1) + 2\delta$  for some  $\delta > 0$ . Then the above obtained inequality  $2t_0 - t_1 \leq f(x_1)$  entails that  $t_1 - \delta \geq t_0$ . Thus

$$\rho|t_1 - \delta - t_0| + \|x_1 - x_0\|^2 = \rho(t_1 - \delta - t_0) + \|x_1 - x_0\|^2 = -\rho\delta + \rho^2 L^2 < \rho^2 L^2,$$

and hence  $\|(x_1, t_1 - \delta) - (x_0, t_0)\| < \rho$ , which is a contradiction since  $(x_1, t_1 - \delta) \in \text{epi } f$ . Thus (4) reads as

$$\rho(f(x_1) - t_0) + \|x_1 - x_0\|^2 = \rho^2 L^2. \quad (5)$$

*Step 3.* In this step we show that if  $x_1$  is such that  $(x_1, f(x_1))$  is a nearest point in  $\text{epi } f$  to  $(x_0, t_0)$  with distance  $\rho$ , then  $x_1$  is a minimum point for the function  $f + \frac{1}{\rho}\|\cdot - x_0\|^2$ . To this end, first fix any  $x \in X$  satisfying  $f(x) \geq t_0$ . We then have  $\|(x, f(x)) - (x_0, t_0)\| \geq \rho$ , and (1) with (5) yield

$$\rho(f(x) - t_0) + \|x - x_0\|^2 \geq \rho^2 L^2 = \rho(f(x_1) - t_0) + \|x_1 - x_0\|^2.$$

Therefore

$$f(x) + \frac{1}{\rho}\|x - x_0\|^2 \geq f(x_1) + \frac{1}{\rho}\|x_1 - x_0\|^2 \quad \text{whenever } x \in X \text{ and } f(x) \geq t_0. \quad (6)$$

Now, fix for a while any  $y \in X$ , with  $f(y) < t_0$ . Then  $(y, t_0) \in \text{epi } f$ , and so  $\|(y, t_0) - (x_0, t_0)\| \geq \rho$ , that is,  $\rho|t_0 - t_0| + \|y - x_0\|^2 \geq \rho^2 L^2$ , which yields  $\|y - x_0\| \geq \rho L$ . Put  $x_2 = x_0 + \frac{\rho L}{\|y - x_0\|}(y - x_0)$ ; then  $\|x_2 - x_0\| = \rho L$ , and we claim that  $f(x_2) \geq t_0$ . Indeed, assume  $f(x_2) < t_0$ . From the continuity of  $f$  there is  $x_3$  in the open linear segment, with the end points  $x_2$  and  $x_0$ , such that  $f(x_3) < t_0$ . Thus  $(x_3, t_0) \in \text{epi } f$  and  $\|x_3 - x_0\| < \|x_2 - x_0\| = \rho L$ . It then follows  $\|(x_3, t_0) - (x_0, t_0)\| \geq \rho$ , and so  $\|x_3 - x_0\|^2 = \rho|t_0 - t_0| + \|x_3 - x_0\|^2 \geq \rho^2 L^2$ , a contradiction. Therefore  $f(x_2) \geq t_0$ .

Now, the Lipschitz property of  $f$  and (6), used for  $x := x_2$ , yield

$$\begin{aligned} f(y) &\geq f(x_2) - L\|y - x_2\| \geq f(x_1) - \frac{1}{\rho}\|x_2 - x_0\|^2 + \frac{1}{\rho}\|x_1 - x_0\|^2 - L\|y - x_2\| \\ &= f(x_1) - L\|y - x_0\| + \frac{1}{\rho}\|x_1 - x_0\|^2 \geq f(x_1) - \frac{1}{\rho}\|y - x_0\|^2 + \frac{1}{\rho}\|x_1 - x_0\|^2 \end{aligned}$$

for every  $y \in X$  with  $f(y) < t_0$ . Here we first used that

$$\|y - x_2\| = \|y - x_0\| - \|x_2 - x_0\| = \|y - x_0\| - \rho L$$

and then that  $L \leq \frac{1}{\rho}\|y - x_0\|$ . We have thus proved that the function  $f + c\|\cdot - x_0\|^2$ , with  $c := \frac{1}{\rho}$ , attains its infimum at the point  $x_1$ .

Step 4. Next, we shall show that the problem of minimizing the function  $f + c\|\cdot - x_0\|^2$  is well-posed. To this end, assume that a sequence  $(y_n)_{n \in \mathbb{N}}$  in  $X$  is such that

$$f(y_n) + \frac{1}{\rho}\|y_n - x_0\|^2 \longrightarrow \inf \left( f + \frac{1}{\rho}\|\cdot - x_0\|^2 \right) \left( = f(x_1) + \frac{1}{\rho}\|x_1 - x_0\|^2 \right) \text{ as } n \rightarrow \infty. \tag{7}$$

From (7) we can easily deduce that  $(y_n)_{n \in \mathbb{N}}$  and  $(f(y_n))_{n \in \mathbb{N}}$  are bounded sequences. We claim that  $\liminf_{n \rightarrow \infty} f(y_n) \geq t_0$ . Indeed, assume, by contradiction, that there are  $\delta > 0$  and an infinite set  $N \subset \mathbb{N}$  such that  $f(y_n) < t_0 - \delta$  for every  $n \in N$ . Find  $\Delta > \frac{\delta}{L}$  so big that  $\|y_n - x_0\| < \Delta$  for every  $n \in N$ . Put

$$z_n = \frac{\delta}{L\Delta}x_0 + \left(1 - \frac{\delta}{L\Delta}\right)y_n, \quad n \in N.$$

Fix any  $n \in N$ . Then

$$\begin{aligned} f(z_n) &\leq f(y_n) + L\|z_n - y_n\| < t_0 - \delta + L\|z_n - y_n\| \\ &= t_0 - \delta + L \cdot \frac{\delta}{L\Delta}\|x_0 - y_n\| \leq t_0, \end{aligned}$$

and hence  $(z_n, t_0) \in \text{epi } f$ . Thus  $\|(z_n, t_0) - (x_0, t_0)\| \geq \rho$ , and so  $\rho|t_0 - t_0| + \|z_n - x_0\|^2 \geq \rho^2 L^2$ , that is,  $\|z_n - x_0\| \geq \rho L$ . But then

$$\|y_n - x_0\| = \frac{1}{1 - \frac{\delta}{L\Delta}}\|z_n - x_0\| \geq \frac{\rho L^2 \Delta}{L\Delta - \delta} > 0 \text{ for every } n \in N.$$

Now, the Lipschitz property of  $f$  and the latter inequality used twice yield

$$\begin{aligned} f(y_n) + \frac{1}{\rho}\|y_n - x_0\|^2 &\geq f(x_0) - L\|y_n - x_0\| + \frac{1}{\rho}\|y_n - x_0\|^2 \\ &\geq f(x_0) - L\|y_n - x_0\| + \frac{L^2 \Delta}{L\Delta - \delta}\|y_n - x_0\| \\ &= f(x_0) + \frac{\delta L}{L\Delta - \delta}\|y_n - x_0\| \geq f(x_0) + \frac{\delta L}{L\Delta - \delta} \cdot \frac{\rho L^2 \Delta}{L\Delta - \delta}. \end{aligned}$$

Therefore,

$$\begin{aligned} (0 <) \frac{\rho \delta L^3 \Delta}{(L\Delta - \delta)^2} &\leq -f(x_0) + \lim_{n \in N} \left( f(y_n) + \frac{1}{\rho}\|y_n - x_0\|^2 \right) \\ &= -f(x_0) + \inf \left( f + \frac{1}{\rho}\|\cdot - x_0\|^2 \right) (\leq 0), \end{aligned}$$

a contradiction. We thus have proved that  $\liminf_{n \rightarrow \infty} f(y_n) \geq t_0$ .

Step 5. We shall show that

$$\rho_n := \|(y_n, f(y_n)) - (x_0, t_0)\| \longrightarrow \rho \text{ as } n \rightarrow \infty. \tag{8}$$

In order to prove (8), it is enough to show that every subsequence of  $(\rho_n)$  has a further subsequence which is convergent to  $\rho$ . So let  $(\rho_{n_k})_{k \in \mathbb{N}}$  be an arbitrary subsequence of

$(\rho_n)$ . Since, as we noticed above, the sequences  $(y_n)_{n \in \mathbb{N}}$  and  $(f(y_n))_{n \in \mathbb{N}}$  are bounded, so is  $(\rho_n)_{n \in \mathbb{N}}$ , and therefore, there are further subsequences of  $(\rho_{n_k})$ ,  $(f(y_{n_k}))$  and  $(y_{n_k})$  (which, for simplicity we will index again by  $n_k, k \in \mathbb{N}$ ) such that  $\lim_k \rho_{n_k}$ ,  $\lim_k f(y_{n_k})$  and  $\lim_k \|y_{n_k} - x_0\|$  exist. Thus, using (7) and (5) we have

$$\begin{aligned} \rho \left( \lim_k f(y_{n_k}) - t_0 \right) + \lim_k \|y_{n_k} - x_0\|^2 &= \rho \lim_k \left( f(y_{n_k}) + \frac{1}{\rho} \|y_{n_k} - x_0\|^2 \right) - \rho t_0 \\ &= \rho \left( f(x_1) + \frac{1}{\rho} \|x_1 - x_0\|^2 \right) - \rho t_0 \\ &= \rho (f(x_1) - t_0) + \|x_1 - x_0\|^2 = \rho^2 L^2. \end{aligned} \tag{9}$$

From the definition of the norm  $\|\cdot\|$  we have

$$\rho_n |f(y_n) - t_0| + \|y_n - x_0\|^2 = \rho_n^2 L^2 \quad \text{for all } n = 1, 2, 3, \dots,$$

and hence, as according to Step 4,  $\lim_k f(y_{n_k}) \geq t_0$ , we have

$$\lim_k \rho_{n_k} \left( \lim_k f(y_{n_k}) - t_0 \right) + \lim_k \|y_{n_k} - x_0\|^2 = \lim_k \rho_{n_k}^2 L^2. \tag{10}$$

And, by subtracting (10) from (9), we get

$$\left( \rho - \lim_k \rho_{n_k} \right) \left( \lim_k f(y_{n_k}) - t_0 \right) = \left( \rho^2 - \lim_k \rho_{n_k}^2 \right) L^2.$$

If  $\lim_k \rho_{n_k} \neq \rho$ , then we get  $\lim_k f(y_{n_k}) - t_0 = (\rho + \lim_k \rho_{n_k}) L^2$ , and regarding (10), we have

$$\begin{aligned} \lim_k \rho_{n_k} \left( \rho + \lim_k \rho_{n_k} \right) L^2 + \lim_k \|y_{n_k} - x_0\|^2 &= \lim_k \rho_{n_k}^2 L^2, \\ \lim_k \rho_{n_k} \rho L^2 + \lim_k \|y_{n_k} - x_0\|^2 &= 0. \end{aligned}$$

And this is impossible since  $\rho > 0$  and  $\rho_n \geq \rho$  for all  $n = 1, 2, 3, \dots$ . Therefore,  $\lim_k \rho_{n_k} = \rho$ . Since we have started with an arbitrary subsequence of  $(\rho_n)$  this entails that  $\lim_{n \rightarrow \infty} \rho_n = \rho$ . We thus have proved (8).

Now, since (8) means that  $(y_n, f(y_n))_{n \in \mathbb{N}}$  is a minimizing sequence for the best approximation problem of  $(x_0, t_0)$  to  $\text{epi } f$ , using *Step 2*, we get that the sequence  $((y_n, f(y_n))_{n \in \mathbb{N}}$  has a norm-cluster point. Hence so does the minimizing sequence  $(y_n)_{n \in \mathbb{N}}$ . (Since the norm  $\|\cdot\|$  is not assumed to be strictly convex, there is no guarantee that a cluster point of  $(y_n)_{n \in \mathbb{N}}$  must be equal to  $x_1$ ; see also Remark 2.2 below). We have thus proved that the problem of minimizing the function  $f + \frac{1}{\rho} \|\cdot - x_0\|^2$  is well-posed.

*Step 6.* Keeping still the same fixed  $x_0$ , it remains to show the residuality of the  $c$ 's in question (so far we have found only one  $c$ ). Consider the function  $\varphi : (-\infty, f(x_0)) \rightarrow (0, +\infty)$  defined by

$$\varphi(\tau) = [\text{dist}((x_0, \tau), \text{epi } f)]^{-1}, \quad \tau \in (-\infty, f(x_0)).$$

It is easy to verify that  $\varphi$  is locally Lipschitzian, and that  $\varphi(\tau) \rightarrow +\infty$  as  $\tau \uparrow f(x_0)$ . Also, the Lipschitz property of  $f$  easily yields that  $\varphi(\tau) \rightarrow 0$  as  $\tau \downarrow -\infty$ . We shall show that  $\varphi$  is strictly increasing. So fix any  $\tau_1 < \tau_2 < f(x_0)$ . Find  $\tau \in (\tau_1, \tau_2)$  so that  $\tau \in S$

( $S$  is the residual subset of  $(-\infty, f(x_0))$  from Step 2). Then there is  $(u, t) \in \text{epi } f$  so that  $\varphi(\tau) = \|\|(x_0, \tau) - (u, t)\|^{-1}$ . We have  $(u, t + \tau_2 - \tau) \in \text{epi } f$ , and this point actually lies in the interior of  $\text{epi } f$  (as  $f$  is continuous). Thus

$$\text{dist}((x_0, \tau_2), \text{epi } f) < \|\|(x_0, \tau_2) - (u, t + \tau_2 - \tau)\| = \|\|(x_0, \tau) - (u, t)\|.$$

Hence  $\varphi(\tau_2) > \varphi(\tau)$ . Now, consider any  $\varepsilon > 0$  and find  $(v, \sigma) \in \text{epi } f$  so that

$$\text{dist}((x_0, \tau_1), \text{epi } f) + \varepsilon > \|\|(x_0, \tau_1) - (v, \sigma)\|.$$

Then  $(v, \sigma + \tau - \tau_1) \in \text{epi } f$ , and so

$$\begin{aligned} \varphi(\tau)^{-1} &= \text{dist}((x_0, \tau), \text{epi } f) \leq \|\|(x_0, \tau) - (v, \sigma + \tau - \tau_1)\| \\ &= \|\|(x_0, \tau_1) - (v, \sigma)\| < \varphi(\tau_1)^{-1} + \varepsilon. \end{aligned}$$

Hence, letting  $\varepsilon \downarrow 0$ , we get  $\varphi(\tau) \geq \varphi(\tau_1)$ . Therefore  $\varphi(\tau_2) > \varphi(\tau_1)$ .

Putting together the above facts we get that  $\varphi$  is a homeomorphism from  $(-\infty, f(x_0))$  onto  $(0, +\infty)$ . Hence, as  $S$  is a residual subset of  $(-\infty, f(x_0))$ , the set  $R_{x_0} := \varphi(S)$  is residual in  $(0, +\infty)$ . And from Step 5 we already know that for every  $c \in R_{x_0}$  the problem of minimizing the function  $f + c\|\cdot - x_0\|^2$  is well-posed. We have thus fully proved (ii).  $\square$

(ii)  $\Rightarrow$  (iii). Assume (ii) holds. Fix any  $z \in X$  and any  $\varepsilon > 0$ . Let  $L > 0$  be a Lipschitzian constant of  $f$ . Let  $\|\|\cdot\|$  be the Minkowski functional of the body  $B$  defined by (1) in Step 1 of the proof of the implication (i)  $\Rightarrow$  (ii); we already know that it is an equivalent norm on  $X \times \mathbb{R}$ . According to (ii) there is  $x_0$  with  $\|x_0 - z\| < \varepsilon/2$  and  $c \in (\frac{2L}{2L+\varepsilon}, 1)$  so that  $f + c\|\cdot - x_0\|^2$  attains its infimum at, say,  $x_1 \in X$ . Thus

$$f(x) + c\|x - x_0\|^2 \geq f(x_1) + c\|x_1 - x_0\|^2 \quad \text{for every } x \in X. \quad (11)$$

The choice  $x := x_0$  and the Lipschitz property of  $f$  yield that  $\|x_1 - x_0\| \leq \frac{L}{c}$ . Put  $t_0 = f(x_1) + c\|x_1 - x_0\|^2 - \frac{L^2}{c}$ . Then  $t_0 \leq f(x_1)$  and

$$\frac{1}{c}(f(x_1) - t_0) + \|x_1 - x_0\|^2 = \frac{1}{c^2}L^2,$$

which means that  $\|\|(x_1, f(x_1)) - (x_0, t_0)\| = \frac{1}{c}$ . Moreover, for  $(x, t) \in \text{epi } f$ , with  $t \geq t_0$ , we have

$$\begin{aligned} \frac{1}{c}(t - t_0) + \|x - x_0\|^2 &\geq \frac{1}{c}(f(x) + c\|x - x_0\|^2 - t_0) \\ &\geq \frac{1}{c}(f(x_1) + c\|x_1 - x_0\|^2 - t_0) = \frac{1}{c^2}L^2, \end{aligned}$$

and so  $\|\|(x, t) - (x_0, t_0)\| \geq \frac{1}{c}$ .

On the other hand, take  $(x, t) \in \text{epi } f$ , with  $t < t_0$ . Then  $f(x) \leq t < t_0$  and, using the definition of  $t_0$  and (11), we obtain

$$t_0 + c\|x - x_0\|^2 > f(x_1) + c\|x_1 - x_0\|^2 = t_0 + \frac{1}{c^2}L^2.$$

Thus  $\|x - x_0\|^2 > \frac{L^2}{c^2}$ , and consequently

$$\frac{1}{c}|t - t_0| + \|x - x_0\|^2 > \frac{L^2}{c^2},$$

showing again that  $\|(x, t) - (x_0, t_0)\| > \frac{1}{c}$ . Therefore, we have shown that the point  $(x_1, f(x_1))$  is a nearest point to  $(x_0, t_0)$  in  $\text{epi } f$  with distance  $\frac{1}{c}$  ( $> 1$ ).

Now, put

$$u = x_1 + c(x_0 - x_1), \quad \tau = f(x_1) + c(t_0 - f(x_1)).$$

We have  $\|(x_1, f(x_1)) - (u, \tau)\| = c\|(x_1, f(x_1)) - (x_0, t_0)\| = 1$  and the point  $(u, \tau)$  lies in the interior of the line segment joining  $(x_0, t_0)$  and  $(x_1, f(x_1))$ . Since  $(x_1, f(x_1))$  is a nearest point to  $(x_0, t_0)$  in  $\text{epi } f$  this means that  $(x_1, f(x_1))$  is a nearest point also to  $(u, \tau)$  with  $\rho = \text{dist}((u, \tau), \text{epi } f) = 1$ . But we already showed in *Step 3* that in such a case the function  $f + \frac{1}{\rho}\|\cdot - u\|^2 = f + \|\cdot - u\|^2$  attains its infimum at  $x_1$ .

Finally, note that

$$\begin{aligned} \|u - z\| &\leq \|u - x_0\| + \|x_0 - z\| = (1 - c)\|x_0 - x_1\| + \|x_0 - z\| \\ &< (1 - c) \cdot \frac{L}{c} + \frac{\varepsilon}{2} = \left(\frac{1}{c} - 1\right)L + \frac{\varepsilon}{2} < \left(\frac{2L + \varepsilon}{2L} - 1\right)L + \frac{\varepsilon}{2} = \varepsilon. \end{aligned}$$

Thus we have completely proved (iii). □

(iii)  $\Rightarrow$  (i). Assume that (iii) holds. As regards the reflexivity of  $X$ , fix any  $\xi \in X^*$ . From (iii) find  $x_0 \in X$ , and  $x_1 \in X$  so that  $\xi(x) + \|x - x_0\|^2 \geq \xi(x_1) + \|x_1 - x_0\|^2$  for all  $x \in X$ . Rearranging this inequality we get

$$\xi(x - x_1) \geq \|x_1 - x_0\|^2 - \|x - x_0\|^2 \quad \text{for all } x \in X. \tag{12}$$

Assume that  $x_1 = x_0$ . Then  $\xi(x - x_0) \geq 0$  for every  $x \in X$  and hence  $\xi = 0$ . Further assume  $x_1 \neq x_0$ . Consider any  $t \in (0, 1)$ . Taking  $x := x_0 + t(x_1 - x_0)$  in (12), we have

$$\begin{aligned} (1 - t)\xi(x_0 - x_1) &= \xi(x_0 + t(x_1 - x_0) - x_1) \\ &\geq \|x_1 - x_0\|^2 - \|x_0 + t(x_1 - x_0) - x_0\|^2 = (1 - t^2)\|x_1 - x_0\|^2. \end{aligned}$$

Dividing both sides here by  $(1 - t)$ , we get  $\xi(x_0 - x_1) \geq (1 + t)\|x_1 - x_0\|^2$ . And letting  $t \uparrow 1$ , we obtain

$$\xi(x_0 - x_1) \geq 2\|x_1 - x_0\|^2. \tag{13}$$

On the other hand, fixing any  $h \in X$ , with  $\|h\| = 1$ , and any  $t > 0$ , when putting  $x := x_1 - th$  in (12), we get

$$\begin{aligned} -t\xi(h) &= \xi(x_1 - th - x_1) \geq \|x_1 - x_0\|^2 - \|x_1 - th - x_0\|^2 \\ &\geq \|x_1 - x_0\|^2 - \|x_1 - x_0\|^2 - 2t\|x_1 - x_0\| - t^2 = -t(2\|x_1 - x_0\| + t). \end{aligned}$$

Here, keeping still  $h$  fixed, then dividing by  $(-t)$ , and finally letting  $t \downarrow 0$ , we get  $\xi(h) \leq 2\|x_1 - x_0\|$ . Thus  $\|\xi\| \leq 2\|x_1 - x_0\|$ . Therefore, comparing this inequality with (13), we get

$$2\|x_1 - x_0\|^2 \leq \xi(x_0 - x_1) \leq \|\xi\|\|x_1 - x_0\| \leq 2\|x_1 - x_0\|^2,$$



and so, since  $x_1 \neq x_0$ , we get

$$\xi \left( \frac{x_0 - x_1}{\|x_1 - x_0\|} \right) = 2\|x_1 - x_0\| = \|\xi\|.$$

Thus, in both cases, our  $\xi$  attains its norm. Then James' theorem guarantees the reflexivity of  $X$ .

**Remark 1.3.** As mentioned by the referee, the fact that  $\xi$  attains its norm, provided (iii) is true, could be derived in a shorter way, using the sum rule for subdifferentials of convex functions (applied for the functions  $\xi$  and  $\|\cdot - x_0\|^2$  above). We have preferred to give here the direct proof of this fact.

Now we shall check that the norm  $\|\cdot\|$  is Kadec-Klee. Assume (iii) holds and that  $\|\cdot\|$  is not Kadec-Klee. We shall profit from a counterexample due to Konjagin [11], [2]. Using [2, Lemma 5.9], we find  $\delta \in (0, \frac{1}{4})$ ,  $\xi \in X^*$ , and  $y_n \in 2B_X \cap \xi^{-1}(1)$ ,  $n \in \mathbb{N}$ , such that  $\lim_{n \rightarrow \infty} \|y_n\| = 1 = \|\xi\|$  and  $\|y_n - y_m\| > 9\delta$  whenever  $n, m \in \mathbb{N}$  are distinct. Find  $k \in \mathbb{N}$  such that  $|\|y_n\| - 1| < \frac{\delta}{4}$  and  $2^{-n} < \frac{\delta}{4}$  for every  $n \geq k$ . Put

$$M = \bigcup_{n=k}^{\infty} M_n \quad \text{where } M_n = (1 + 2^{-n})y_n + (3\delta B_X \cap \xi^{-1}(0)).$$

Here, each  $M_n$  is closed, and for all distinct  $n, m \in \{k, k+1, \dots\}$  the distance between  $M_n$  and  $M_m$  is at least  $\delta$ . Therefore the whole set  $M$  is closed. Take  $\alpha > \max\{4, \frac{4}{\delta}(1 + \delta)^2\}$ . Define then the function  $f : X \rightarrow [0, +\infty)$  by  $f = \alpha \inf\{\|\cdot - u\|; u \in M\}$ ; note that  $f$  is Lipschitzian. We shall show that for any  $x_0 \in \frac{\delta}{4}B_X$  the function  $f + \|\cdot - x_0\|^2$  does not attain its infimum. So fix such an  $x_0$  and assume, by contradiction, that there is  $x_1 \in X$  so that

$$f(x) + \|x - x_0\|^2 \geq f(x_1) + \|x_1 - x_0\|^2 \quad \text{for all } x \in X. \tag{14}$$

We shall show that  $x_1$  must lie in  $M$ . Put  $u = y_k(1 + 2^{-k})$ . Then  $u \in M_k \subset M$ , and so

$$\begin{aligned} f(u) + \|u - x_0\|^2 &= \|y_k + 2^{-k}y_k - x_0\|^2 \leq \left( \|y_k\| + \frac{\delta}{4}\|y_k\| + \|x_0\| \right)^2 \\ &< \left( 1 + \frac{\delta}{4} + \frac{\delta}{4} \left( 1 + \frac{\delta}{4} \right) + \frac{\delta}{4} \right)^2 < (1 + \delta)^2. \end{aligned}$$

Now, we claim that  $\frac{1}{\alpha}f(x_1) < \frac{\delta}{4}$ . Indeed, otherwise, using the latter inequality and the choice of  $\alpha$ , we would have

$$f(x_1) + \|x_1 - x_0\|^2 \geq \frac{\alpha\delta}{4} > (1 + \delta)^2 > f(u) + \|u - x_0\|^2,$$

which contradicts to (14). Thus,  $\frac{1}{\alpha}f(x_1) < \frac{\delta}{4}$ , and since the distance between the sets  $M_n$  and  $M_m$  for distinct  $n, m \geq k$  is greater than  $\delta$ , this entails that  $\frac{1}{\alpha}f(x_1) = \inf\{\|x_1 - y\|; y \in M_{n_1}\}$  for some (unique)  $n_1 \geq k$ .

Since each  $M_n$  is closed convex and bounded, and since  $X$  is already shown to be reflexive, there must exist  $y_1 \in M_{n_1}$  such that  $\|x_1 - y_1\| = \frac{1}{\alpha}f(x_1)$ . Assume that  $x_1 \notin M_{n_1}$ . Then,

$\|x_1 - y_1\| > 0$ , and using the fact that  $\alpha > 2\|y_1 - x_0\|$ , we can estimate

$$\begin{aligned} & f(x_1) + \|x_1 - x_0\|^2 \\ &= \alpha\|x_1 - y_1\| + \|x_1 - x_0\|^2 \geq \alpha\|x_1 - y_1\| + (\|y_1 - x_0\| - \|x_1 - y_1\|)^2 \\ &= \alpha\|x_1 - y_1\| + (\|y_1 - x_0\|^2 - 2\|y_1 - x_0\|\|x_1 - y_1\| + \|x_1 - y_1\|^2) \\ &> (\alpha - 2\|y_1 - x_0\|)\|x_1 - y_1\| + \|y_1 - x_0\|^2 > \|y_1 - x_0\|^2 = f(y_1) + \|y_1 - x_0\|^2, \end{aligned}$$

which contradicts (14). Therefore  $x_1 \in M_{n_1} \subset M$ .

Now, for every  $n \geq k$  and every  $x \in M_n$  we thus have from (14) that  $\|x - x_0\|^2 \geq \|x_1 - x_0\|^2$  and hence  $\|x - x_0\| \geq \|x_1 - x_0\|$ . Fix any  $n \geq k$ . Put  $w_n = (1 + 2^{-n})y_n + x_0 - \xi(x_0)y_n$ . Then  $\|w_n - (1 + 2^{-n})y_n\| \leq \|x_0\| + 2\|x_0\| \leq 3\delta$  and  $\xi(w_n - (1 + 2^{-n})y_n) = 0$ . Hence  $w_n \in M_n$  and we thus have  $\|w_n - x_0\| \geq \|x_1 - x_0\|$ . Moreover,

$$\|w_n - x_0\| = (1 + 2^{-n} - \xi(x_0))\|y_n\| \rightarrow 1 - \xi(x_0) \text{ as } n \rightarrow \infty.$$

Therefore

$$1 - \xi(x_0) = \lim_{n \rightarrow \infty} \|w_n - x_0\| \geq \|x_1 - x_0\| \geq \xi(x_1 - x_0) = 1 + 2^{-n_1} - \xi(x_0),$$

a contradiction. We have completely proved (i). □

(i)  $\Rightarrow$  (ii'). Assume that the function  $f : X \rightarrow (-\infty, +\infty]$  is proper, lower semicontinuous and bounded below. We shall imitate the argument from the proof of the implication (i)  $\Rightarrow$  (ii). Take  $B = \{(x, t) \in X \times \mathbb{R}; |t| \leq -\|x\|^2 + 1\}$  and let  $\|\cdot\|$  be the Minkowski functional of  $B$ . Let the sets  $O, E, \Omega$ , and  $x_0 \in \Omega$  be found for this norm  $\|\cdot\|$  as before. We find a number  $t_0 < \inf f$  so that  $(x_0, t_0) \in E$ . Let  $(x_1, t_1) \in \text{epi } f$  and  $\rho$  be found for our  $(x_0, t_0)$  so that (2) is satisfied. Then for sure  $t_1 > t_0$  and the formula (3) has the form

$$\rho(t_1 - t_0) + \|x_1 - x_0\|^2 = \rho^2.$$

And, as in the proof of the implication (i)  $\Rightarrow$  (ii), we can see that actually  $t_1 = f(x_1)$ . Thus we have formula (5) with  $L = 1$ .

Now, for any  $x \in X$  we have  $f(x) > t_0$ . Hence the reasoning from the proof of (i)  $\Rightarrow$  (ii) leading to formula (6) in *Step 3* yields

$$f(x) + \frac{1}{\rho}\|x - x_0\|^2 \geq f(x_1) + \frac{1}{\rho}\|x_1 - x_0\|^2 \text{ whenever } x \in X.$$

Further we proceed as in the proof of (i)  $\Rightarrow$  (ii). The claim in *Step 4* there is satisfied automatically. *Step 5* works without any change. Then the problem of minimizing the function  $f + \frac{1}{\rho}\|\cdot - x_0\|^2$  is well-posed.

It remains to prove *Step 6*. For the still fixed  $x_0$  we put  $a_{x_0} = \varphi(\inf f)$  (the latter could be  $\infty$ ), where the function  $\varphi$  is as in *Step 6*. One easily sees that  $\varphi(\tau) \rightarrow a_{x_0}$  as  $\tau \uparrow \inf f$ . The fact that  $f$  is bounded below yields that  $\varphi(\tau) \rightarrow 0$  as  $\tau \downarrow -\infty$ . We need to show that  $\varphi$  is strictly increasing. As in *Step 6* above, fix  $\tau_1 < \tau_2 < \inf f$  and take  $\tau \in (\tau_1, \tau_2)$  so that  $\tau \in S$  ( $S$  is the residual subset of  $(-\infty, f(x_0))$  from *Step 2* corresponding to  $x_0$ ).

Then there is  $(u, t) \in \text{epi } f$  so that  $\varphi(\tau) = \|(x_0, \tau) - (u, t)\|^{-1} =: \frac{1}{\rho_0}$ . Thus, according to the definition of the norm  $\|\cdot\|$ , we have (since  $t \geq \inf f$ )

$$\rho_0(t - \tau) + \|x_0 - u\|^2 = \rho_0^2.$$

On the other hand, putting  $\rho = \|(x_0, \tau_2) - (u, t)\|$ , then

$$\rho(t - \tau_2) + \|x_0 - u\|^2 = \rho^2.$$

Consequently

$$\rho_0^2 - \rho_0(t - \tau) = \rho^2 - \rho(t - \tau_2) = \|x_0 - u\|^2 (\geq 0).$$

And, since  $t - \tau > t - \tau_2 > 0$ , we conclude that  $\rho < \rho_0$ . Therefore

$$\text{dist}((x_0, \tau_2), \text{epi } f) \leq \rho < \rho_0,$$

which gives  $\varphi(\tau_2) > \varphi(\tau)$ . The fact that  $\varphi(\tau) \geq \varphi(\tau_1)$  is proved as in *Step 6*.

Then clearly there is a residual set  $R_{x_0} \subset (0, a_{x_0})$  satisfying *(ii')* for  $x_0$ . We have thus verified *(ii')*.  $\square$

*(ii')*  $\Rightarrow$  *(iii')* is trivial.

*(iii')*  $\Rightarrow$  *(i)*. Suppose *(iii')* is true. In order to prove the reflexivity of  $X$  let us fix any  $\xi \in X^*$ , with  $\|\xi\| = 1$ . Define  $f : X \rightarrow \mathbb{R}$  by

$$f(x) = \begin{cases} 0 & \text{if } x \in \xi^{-1}(0), \\ +\infty & \text{if } x \in X \setminus \xi^{-1}(0). \end{cases}$$

Clearly,  $f$  is lower semicontinuous and bounded below. To this  $f$ , let  $D \subset X$  be the dense set given by *(iii')*. Pick then  $x_0 \in D \cap (X \setminus \xi^{-1}(0))$  and let  $c$  be such that  $f + c\|\cdot - x_0\|^2$  attains its minimum at some  $x_1$ , i.e.  $f + c\|\cdot - x_0\|^2 \geq f(x_1) + c\|x_1 - x_0\|^2$ . We necessarily have that  $\xi(x_1) = 0$ . Thus, for every  $x \in \xi^{-1}(0)$  we have  $c\|x - x_0\|^2 \geq c\|x_1 - x_0\|^2$ , that is,  $\|x - x_0\| \geq \|x_1 - x_0\| (> 0)$ . We shall show that  $|\xi(x_1 - x_0)| = \|x_1 - x_0\|$ . Assume not; then  $|\xi(x_1 - x_0)| < \|x_1 - x_0\|$ . Find  $z \in X$  so that  $\|z - x_0\| = \|x_1 - x_0\|$  and  $\xi(z - x_0) > \frac{1}{2}(\|x_1 - x_0\| + |\xi(x_1 - x_0)|)$ . Then, putting

$$w = x_0 + \frac{\xi(x_1 - x_0)}{\xi(z - x_0)}(z - x_0),$$

we have  $w \in \xi^{-1}(0)$  and

$$\begin{aligned} \|x_1 - x_0\| &\leq \|w - x_0\| = \frac{|\xi(x_1 - x_0)|}{\xi(z - x_0)} \|z - x_0\| \\ &< \frac{\|x_1 - x_0\|}{\xi(z - x_0)} (2\xi(z - x_0) - \|x_1 - x_0\|) \leq \|x_1 - x_0\|; \end{aligned}$$

a contradiction. Therefore  $|\xi(x_1 - x_0)| = \|x_1 - x_0\|$ , and James' theorem guarantees that  $X$  must be reflexive.

If the norm in  $X$  is not Kadec-Klee, then (iii') is violated. Indeed, assume that the norm in  $X$  is not Kadec-Klee and take the closed set  $M$  from the proof of the implication (iii)  $\Rightarrow$  (i). Let the function  $f$  be defined as follows:

$$f(x) = \begin{cases} 0 & \text{if } x \in M, \\ +\infty & \text{if } x \in X \setminus M. \end{cases}$$

Evidently,  $f$  is lower semicontinuous and bounded below. Let  $\delta > 0$  be the number associated with the set  $M$ . We shall show that for any  $x_0 \in \frac{\delta}{4}B_X$  and any  $c > 0$  the function  $f + c\|\cdot - x_0\|^2$  does not attain its infimum. Suppose this is not the case. Then there are  $x_0 \in \frac{\delta}{4}B_X$ ,  $c > 0$ , and  $x_1 \in X$  so that  $f(x) + c\|x - x_0\|^2 \geq f(x_1) + c\|x_1 - x_0\|^2$  for every  $x \in X$ . Clearly  $x_1 \in M$ , and so we get  $\|x - x_0\| \geq \|x_1 - x_0\|$  for every  $x \in M$ . Now we proceed exactly as at the end of the the proof of the implication (iii)  $\Rightarrow$  (i) to get a contradiction. We have thus completely verified (i), which ends the proof of Theorem 1.1.  $\square$

The genericity of the set of perturbations yields the following

**Corollary 1.4.** *Let  $(X, \|\cdot\|)$  be a reflexive Banach space, with Kadec-Klee norm, and let a function  $f : X \rightarrow \mathbb{R}$  be either Lipschitzian or be continuous and bounded. Then there exists a residual set  $\Omega \subset X$  such that for every  $u \in \Omega$  there are  $a_u \in (0, +\infty]$  and a residual set  $R_u \subset (0, a_u)$  so that for each  $c \in R_u$   $\inf (f + c\|\cdot - u\|^2)$  and  $\sup (f - c\|\cdot - u\|^2)$  are both attained. (In the Lipschitzian case  $a_u = +\infty$ .)*

## 2. Concluding remarks and results

This section contains several remarks strengthening some of the conclusions in our principle as well as giving some corollaries. We also give some relations to other similar variational principles.

**Remark 2.1.** To prove our Theorem 1.1 we used the result of K.-S. Lau [13, Theorem 4], asserting that *given a closed nonempty set  $C$  in a reflexive Banach space  $X$  with Kadec-Klee norm, there is a residual subset  $\Omega$  of  $X$  such that for each  $x \in \Omega$  the best approximation problem of  $x$  to  $C$  is well-posed.* Actually, Theorem 1.1 contains this result as a special case. This can be readily seen by taking such a  $C \subset X$  and applying (ii') to the function  $f : X \rightarrow (-\infty, +\infty]$  given by  $f(x) = 0$  if  $x \in C$  and  $f(x) = +\infty$  otherwise. We believe that a direct proof of our theorem using Baire category argument exists without using the Lau's result (see also our Remark 2.6 below).

**Remark 2.2 (Uniqueness of the minimum point for the perturbations).** If the problem to minimize a given proper bounded from below function  $g : X \rightarrow (-\infty, +\infty]$  is well-posed and with unique minimizer, then actually every minimizing sequence for  $g$  converges to this unique minimizer. This property is known in Optimization as *the minimization problem for  $g$  is Tykhonov well-posed*, and in Analysis as  *$g$  admits a strong minimum*. Lau's result [13, Theorem 4] can be used (see [2, Theorem 6.6]) to show that, if the norm in the reflexive space  $X$  is both Kadec-Klee and strictly convex, then for every nonempty closed subset  $C$  of  $X$  the problem of best approximation in  $C$  is generically (in  $X$ ) well-posed with unique solution. Thus, there exists a variant of Theorem 1.1: if in (i) we put *The space  $X$  is reflexive and the norm is strictly convex and Kadec-Klee*, this

will be equivalent to all of the other assertions simply adding in them that the minimizer for the corresponding perturbations is unique. We omit rather routine proofs.

**Remark 2.3.** The norm  $\|\cdot\|$  in a reflexive space  $X$  can be taken, both Kadec-Klee and Fréchet smooth according to well known renormings. Thus we get from Theorem 1.1 a “Fréchet smooth” variational principle. If  $X$  is even superreflexive, then it admits an equivalent norm  $\|\cdot\|$  which is Kadec-Klee, and moreover has a modulus of smoothness of power type, that is, there are  $1 < p \leq 2$  and  $c > 0$  such that  $\|x + th\| + \|x - th\| - 2 \leq ct^p$  whenever  $x, h$  lie in the unit sphere of  $X$ , and  $t > 0$ ; then our principle provides supporting by an even smoother function.

**Remark 2.4.** Given a Lipschitz function  $f$  on  $X$ , the set of points of supporting by a parabola may not be residual. Indeed, if so, then, when applying our principle also for  $-f$ , we would get that  $f$  is Fréchet differentiable at the points of a residual set, contradictory to a known counterexample in Hilbert space due to Preiss [16].

**Remark 2.5.** If  $X$  is reflexive, with Kadec-Klee norm  $\|\cdot\|$ , and  $f : X \rightarrow \mathbb{R}$  is lower semicontinuous with no further properties, then it may happen that  $f + c\|\cdot - u\|^2$  attains its infimum for no  $c \geq 0$  and for no  $u \in X$ . Indeed, define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by  $f(t) = t^3$ ,  $t \in \mathbb{R}$ . Then  $\inf_{t \in \mathbb{R}} (f(t) + c(t - u)^2) = -\infty$  for every  $c \geq 0$  and every  $u \in \mathbb{R}$ .

**Remark 2.6.** Our argument does not guarantee if, in (ii) of Theorem 1.1, we can have  $c = 1$  for some  $u \in \Omega$ . This is done in (iii) for the price of eventual loss of the residuality of the set of  $u$ 's and eventual loss of the well-posedness of the problems of minimizing  $f + \|\cdot - u\|^2$ . In fact, it turns out that in (iii) one can have the residuality of the points giving the perturbing parabolas, as it is shown in the forthcoming paper [7], which contains also a more direct approach to the proof of (iii). This proof is based on a technique of K.-S. Lau [13].

Note that in (iii) we can replace  $f + \|\cdot - u\|^2$  by  $f + c\|\cdot - u\|^2$  with an arbitrary a priori fixed  $c > 0$ , just by considering the norm  $\sqrt{c}\|\cdot\|$ .

**Remark 2.7 (Relations with Borwein-Preiss smooth variational principle).**

This principle is usually understood as that with a perturbing function in form of the sum of countably many parabolas, see [2, Theorem 2.6], [15, Theorems 4.20]. Yet in [2, Theorem 5.2], dealing with reflexive spaces with Kadec norm, a perturbing function is just one parabola. This is very close to (iii') in our Theorem 1.1. In (ii) and (ii') we have in addition well-posedness of the perturbations as well as *genericity* of the points  $u$  which generate perturbing parabolas, while Borwein-Preiss principles do not have it. Moreover, our Theorem 1.1 covers also the case of Lipschitz, not necessarily bounded below, functions. Let us mention that genericity of the perturbations occurs also in the Stegall variational principle [17, 18] (see also below) and Deville-Godefroy-Zizler variational principle [4] (cf. also a strengthening of the latter in [5]).

On the other hand, unlike of Borwein-Preiss principle we do not have the same kind of control over the location of the minimum point. What we have as a direct consequence from formula (5) is that the minimum point  $x_1$  of  $f + c\|\cdot - x_0\|^2$  found in the proofs of (i)  $\Rightarrow$  (ii) (and (i)  $\Rightarrow$  (ii')) satisfies  $\|x_1 - x_0\| < \frac{L}{c}$  (with  $L = 1$  in the non-Lipschitzian case). This means that in the Lipschitzian case we can have a minimum point of the perturbation as close to a given point as we want, at the price of having the coefficient

$c$  of the perturbing parabola big. In particular, this shows that in such a case we have density of the points realizing the minima of the perturbations.

The same kind of control could be obtained also in the case of a proper lower semicontinuous bounded from below function  $f$  which is continuous at the points of its effective domain  $\text{dom } f$  (the latter implies that  $\text{dom } f$  is open). To see this, let  $\varepsilon > 0$ . Then, by the proof of (i)  $\Rightarrow$  (ii') there is a residual set  $\Omega'$  in the nonempty open set  $\{x \in \text{dom } f; f(x) < \inf f + \varepsilon\}$  so that for each  $x_0 \in \Omega'$  there is  $a_{x_0} > 0$  such that  $f + c\|\cdot - x_0\|^2$  attains its minimum for residually many  $c \in (0, a_{x_0})$  at some corresponding  $x_1$ . Formula (5), with  $L = 1$ , gives  $\|x_1 - x_0\| \leq \frac{1}{c}$ . On the other hand, we know from Step 6 that  $a_{x_0} = \lim_{\tau \uparrow \inf f} [\text{dist}((x_0, \tau), \text{epi } f)]^{-1}$ . Since for  $\tau < \inf f$  we have

$$\text{dist}((x_0, \tau), \text{epi } f) \leq \| (x_0, \tau) - (x_0, f(x_0)) \| = f(x_0) - \tau,$$

we obtain that  $a_{x_0} > \frac{1}{\varepsilon}$ . Thus for residually many  $c \in (\frac{1}{\varepsilon}, a_{x_0})$  the minimum point  $x_1$  will satisfy  $\|x_1 - x_0\| \leq \frac{1}{c} < \varepsilon$ . Observe that in this case, by formula (5) with  $L = 1$ , we have also  $f(x_1) - t_0 \leq \frac{1}{c}$  which gives  $f(x_1) < \inf f + \frac{1}{c} < \inf f + \varepsilon$  for  $c$  as above. That is, we have also control over the value of  $f$  at  $x_1$ : The minimum point  $x_1$  of  $f + c\|\cdot - x_0\|^2$  is an  $\varepsilon$ -minimum of  $f$ .

**Remark 2.8 (Relations with Stegall variational principle).** Let us recall some notions: Given a nonempty set  $A$  of a Banach space  $X$  a *slice* in  $A$  is every set of the form

$$\left\{ x \in A : x^*(x) > \sup_{y \in A} x^*(y) - \alpha \right\}$$

for some  $x^* \in X^*$  and  $\alpha > 0$ . The set  $A$  is called *dentable* if it admits slices of arbitrary small diameter. The Banach space  $X$  has the *Radon-Nikodým property* (in short RNP) if every nonempty bounded subset of  $X$  is dentable.

The Stegall variational principle [17, 18] (see also [15]) states the following: *Let  $f : C \rightarrow \mathbb{R} \cup \{+\infty\}$  be a proper bounded from below lower semicontinuous function defined on a closed convex and bounded subset  $C$  of a Banach space  $X$  with RNP. Then the set  $\{x^* \in X^* : f + x^*$  attains its strong minimum on  $C\}$  is a residual subset of  $X^*$ .*

This principle can be used to obtain a perturbing function in the form  $\|\cdot\|^2 + x^*$  where  $x^* \in X^*$ . More precisely, we have the following

**Theorem 2.9.** *Let  $(X, \|\cdot\|)$  be a Banach space with RNP. Let  $f : X \rightarrow (-\infty, +\infty]$  be a proper lower semicontinuous function bounded below by  $a - b\|\cdot\|^p$  where  $a \in \mathbb{R}$ ,  $b > 0$ , and  $1 \leq p < 2$ . Then there exists a residual set  $\Omega \subset X^*$  such that for every  $x^* \in \Omega$  the function  $f + \|\cdot\|^2 + x^*$  attains strongly its infimum. If  $(X, \|\cdot\|)$  is a Hilbert space, then there even exists a residual set  $\Omega \subset X$  such that for every  $x^* \in \Omega$  the function  $f + \|\cdot - x^*\|^2$  attains strongly its infimum.*

**Proof.** We can see that

$$\frac{1}{\|x\|} (f(x) + \|x\|^2) \geq \frac{1}{\|x\|} (a - b\|x\|^p + \|x\|^2) \rightarrow +\infty \text{ as } x \in X \text{ and } \|x\| \rightarrow +\infty.$$

Hence, for every  $n \in \mathbb{N}$  there is  $a_n \in \mathbb{R}$  such that  $f + \|\cdot\|^2 \geq a_n + 2n\|\cdot\|$ . Fix  $n > 1$ . We will show that there is a residual subset  $\Omega_n \subset nB_{X^*}$  (where  $B_{X^*}$  is the unit ball in  $X^*$ ) such that for any  $x^* \in \Omega_n$  the function  $f + \|\cdot\|^2 + x^*$  attains strong minimum on  $X$ .

To this end, we follow the lines of the proof of Corollary 5.22 from [15]. Since we can replace  $f + \|\cdot\|^2$  by  $f + \|\cdot\|^2 - a_n$ , we may suppose that  $a_n = 0$ . Let us note that in such a case for any  $x^* \in nB_{X^*}$  and any  $x \in X$  we have

$$f(x) + \|x\|^2 + x^*(x) \geq 2n\|x\| - n\|x\| = n\|x\|. \tag{15}$$

Let  $r := (1/n)(f(0) + 1)$  and set  $C := rB_X$ . Applying now the Stegall variational principle for the restriction  $(f + \|\cdot\|^2)|_C$  of  $f + \|\cdot\|^2$  on  $C$  we obtain a residual subset  $G$  of  $X^*$  such that for any  $x^* \in G$  the function  $(f + \|\cdot\|^2)|_C + x^*$  attains its strong minimum on  $C$ . Put  $\Omega_n := nB_{X^*} \cap G$ . The latter set is residual in  $nB_{X^*}$ . We will show that, in fact, for any  $x^* \in \Omega_n$  the function  $f + \|\cdot\|^2 + x^*$  attains its strong minimum also in  $X$ , not only in  $C$ .

To prove this, let  $x^* \in \Omega_n$  and let  $x_0 \in C$  be the strong minimum of  $f + \|\cdot\|^2 + x^*$  on  $C$ . Take  $x \in X$  and suppose that  $f(x) + \|x\|^2 + x^*(x) \leq f(x_0) + \|x_0\|^2 + x^*(x_0) = \inf_C(f + \|\cdot\|^2 + x^*) \leq f(0)$ . Thus, by (15) we have  $\|x\| \leq (1/n)f(0)$  and hence  $x \in C$  yielding  $x = x_0$ . Therefore,  $x_0$  is the unique minimum of  $f + \|\cdot\|^2 + x^*$  on  $X$ . Further, let  $(x_k)_{k \in \mathbb{N}}$  be a minimizing sequence for  $f + \|\cdot\|^2 + x^*$ , that is  $f(x_k) + \|x_k\|^2 + x^*(x_k) \rightarrow f(x_0) + \|x_0\|^2 + x^*(x_0)$ . Then for  $k$  large enough we will have  $f(x_k) + \|x_k\|^2 + x^*(x_k) < f(0) + 1$ , and therefore, again from (15) we see that  $x_k$  must be in  $C$  and hence  $x_k \rightarrow x_0$ . Thus,  $f + \|\cdot\|^2 + x^*$  attains also its strong minimum on  $X$  and our assertion is proved.

It suffices now to set  $\Omega = \bigcup_{n=1}^\infty \Omega_n$ . This is a residual set in  $X^*$  and satisfies the conclusion of the theorem. The case of a Hilbert space can be obtained immediately from the geometry of the Hilbertian norm. Indeed, we have  $\|x\|^2 - x^*(x) = \|x - \frac{x^*}{2}\|^2 - \frac{1}{4}\|x^*\|^2$  for every  $x \in X$ . □

For Hilbert spaces, the above result gives a stronger conclusion than our statement (iii) because we do not have the residuality of  $u$ 's in (iii). Further, it should be noted that Stegall principle does not need the Kadec-Klee property of the norm neither the reflexivity of the space. Only the fact that  $X$  possesses RNP is assumed (every reflexive space has RNP, see for instance [15, Chapter 5]). Also, the boundedness below by the function  $a - b\|\cdot\|^p$  is guaranteed by both the Lipschitz property of  $f$  as well as by  $\inf f > -\infty$ .

**Remark 2.10.** We note that in a reflexive space any Kadec-Klee norm is already Kadec, which means that the weak and the norm topologies on the sphere coincide. This is a consequence of a more general fact that *in any Asplund space, bounded weakly sequentially closed sets are weakly closed*. Indeed, consider an Asplund space  $X$  and a bounded weakly sequentially closed set  $M \subset X$ , and let  $x$  be an element of the weak closure of  $M$ . By Kaplanski's theorem [10, p. 129], there is a countable set  $C \subset M$  such that  $x$  belongs to the weak closure of  $C$ . Let  $Y$  be the closed linear span of  $C$ . Since  $X$  is Asplund,  $Y^*$  must be separable. Let  $S$  be a countable norm-dense subset in  $B_{Y^*}$ . Find then a sequence  $(x_n)_{n \in \mathbb{N}}$  in  $C$  such that  $y^*(x_n) \rightarrow y^*(x)$  as  $n \rightarrow \infty$  for every  $y^* \in S$ . Then also  $x^*(x_n) \rightarrow x^*(x)$  as  $n \rightarrow \infty$  for every  $x^* \in X^*$ . And since  $M$  was weakly sequentially closed, we conclude that  $x \in M$ .

Finally, let us mention that it might be possible to use some ideas from [2] in order to extend our principle for non-reflexive spaces.

**Acknowledgements.** We thank J. Duda and R. Deville for initial discussions related to some results of this paper.

## References

- [1] E. Bishop, R. R. Phelps: A proof that every Banach space is subreflexive, *Bull. Amer. Math. Soc.* 67 (1961) 97–98.
- [2] J. Borwein, S. Fitzpatrick, Existence of nearest points in Banach spaces, *Canad. J. Math.* 49 (1989) 702–720.
- [3] J. Borwein, D. Preiss: A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions, *Trans. Amer. Math. Soc.* 303 (1987) 517–527.
- [4] R. Deville, G. Godefroy, V. Zizler: A smooth variational principle with applications to Hamilton–Jacobi equations in infinite dimensions, *J. Funct. Anal.* 111 (1993) 197–212.
- [5] R. Deville, J. P. Revalski: Porosity of ill-posed problems, *Proc. Amer. Math. Soc.* 128 (2000) 1117–1124.
- [6] I. Ekeland: Nonconvex minimization problems, *Bull. Amer. Math. Soc.* 1 (1979) 443–474.
- [7] M. Fabian: On minimum principles, *Acta Polytech.* 20 (1983) 109–118.
- [8] M. Fabian: Infimal convolution in Efimov–Stečkin Banach spaces, *J. Math. Anal. Appl.* 339 (2008) 735–739.
- [9] M. Fabian, C. Finet: On Stegall’s smooth variational principle, *Nonlinear Anal., Theory Methods Appl.* 66A (2007) 565–570.
- [10] M. Fabian, P. Habala, P. Hájek, V. Montesinos, J. Pelant, V. Zizler: *Functional Analysis and Infinite-Dimensional Geometry*, Springer, New York (2001).
- [11] S. V. Konjagin: On approximation of closed sets in Banach spaces and the characterization of strongly convex spaces, *Sov. Math., Dokl.* 21 (1980) 418–422.
- [12] K. Kuratowski: *Topologie. Vol. I*, PWN, Warsaw (1958).
- [13] K. S. Lau: Almost Chebyshev subsets in reflexive Banach spaces, *Indiana Univ. Math. J.* 27 (1978) 791–795.
- [14] R. R. Phelps: Dentability and extreme points in Banach spaces, *J. Funct. Anal.* 16 (1974) 78–90.
- [15] R. R. Phelps: *Convex Functions, Monotone Operators, and Differentiability*, Lecture Notes in Mathematics 1364, Springer, Berlin (1993).
- [16] D. Preiss: Gâteaux differentiable Lipschitz functions need not be Fréchet differentiable on a residual subset, *Suppl. Rend. Circ. Mat. Palermo, II. Ser.* (1982) 217–222.
- [17] Ch. Stegall: Optimization of functions on certain subsets of Banach spaces, *Math. Ann.* 236 (1978) 171–176.
- [18] Ch. Stegall: Optimization and differentiation in Banach spaces, *Linear Algebra Appl.* 84 (1986) 191–211.