

Finitely generated connected locally compact groups

Karl H. Hofmann and Sidney A. Morris

Introduction

HOFMANN and MORRIS [6] proved that a locally compact connected group G has a finite subset generating a dense subgroup if and only if the weight $w(G)$ of G does not exceed \mathfrak{c} , the cardinality of the continuum. The minimum cardinality of such a topological generating set is an invariant of the group, is denoted by $\sigma(G)$, and is called the *topological rank* of G . For compact abelian groups of weight $\leq \mathfrak{c}$, this number is 1. It was also shown there that for any compact connected group G of weight $\leq \mathfrak{c}$, the invariant $\sigma(G) \leq 2$. CLEARY and MORRIS [3] observed that $\sigma(\mathbb{R}^n) = n + 1$, for $n \geq 1$ and proved the surprising result that for any compact connected group G with $w(G) \leq \mathfrak{c}$, the invariant $\sigma(G \times \mathbb{R}^n) = n + 1$ for $n \geq 1$. Here we extend their result significantly. For example, if G is a compact connected group of weight $\leq \mathfrak{c}$ and L is a nonsingleton connected Lie group, then $\sigma(G \times L) = \sigma(L)$. For more general locally compact groups we bound the topological rank of a group in terms of an associated Lie group.

Partial results

We consider here Hausdorff topological groups only. For a subset $X \subseteq G$ we let $\langle X \rangle$ denote the group generated by X .

Lemma 1. *Let P and A denote topological groups, X a topological generating set of A and $\pi: X \rightarrow P$ a function such that $\pi(X)$ is a topological generating set of P . Set $G = P \times A$, $\text{graph}(\pi) = \{(\pi(x), x) : x \in X\}$, and $H = \overline{\langle \text{graph}(\pi) \rangle}$. Let pr_P and pr_A denote the projections. Then*

$$(*) \quad \overline{\text{pr}_P(H)} = P \quad \text{and} \quad \overline{\text{pr}_A(H)} = A.$$

Proof. Since X , resp. $\pi(X)$, is a topological generating set of A , resp., P , then $\text{pr}_P(\pi(X))$, resp. $\text{pr}_A(X)$ topologically generates the group P , resp. A . Thus $\text{pr}_P(H)$ and $\text{pr}_A(H)$ are dense in P and A , respectively. ■

Lemma 2. *Let P and A denote topological groups, and H a closed subgroup of $P \times A$ such that $(*)$ is satisfied. In the notation of Lemma 1:*

- (i) If P is compact, then $\text{pr}_A|_H: H \rightarrow A$ is an open surjective morphism of topological groups.
- (ii) $N \stackrel{\text{def}}{=} ((P \times \{1\}) \cap H) \times ((\{1\} \times A) \cap H)$ is normal in G .
- (iii) Set $N_P = \{p \in P : (p, 1) \in H\}$ and $A_N = \{a \in A : (1, a) \in H\}$. Then G/N can be identified with $P/N_P \times A/N_A$ and H/N with a closed subgroup thereof. Further, $H = G$ iff $N = G$.
- (iv) If $N = \{1\}$ and P is compact, then there is a unique injective morphism $f: A \rightarrow P$ with dense image extending π such that $H = \{(f(a), a) : a \in A\}$.

Proof. (i) Since P is compact, the projection pr_A is a proper map. Hence it maps the closed group H onto its image in such a fashion that the induced map $H/((P \times \{1\}) \cap H) \rightarrow A$ is an isomorphism of topological groups.

(ii) The normalizer of $(P \times \{1\}) \cap H$ contains H (since $P \times \{1\}$ is normal in G) and $\{1\} \times A$ (since $\{1\} \times A$ centralizes $P \times \{1\}$). Since $\text{pr}_P(H)$ is dense in P , then $H(\{1\} \times A)$ is dense in G . Hence $(P \times \{1\}) \cap H$ is normal in G . Likewise $(\{1\} \times A) \cap H$. Thus N is normal.

(iii) is straightforward.

(iv) Since H meets the factor P trivially, H is the graph of a morphism $f: A \rightarrow P$. Since H is closed and P is compact, it is continuous. Since H meets A trivially, f is injective. By (i) the image of f is dense in P . Since $(\pi(x), x) \in H$ for all x , the morphism f extends π . Since X is a topological generating set, the extension is uniquely determined by this condition. ■

Lemma 3. Let P be a compact connected group and A a locally compact group. Assume that H is a closed subgroup of $G = P \times A$ satisfying $(*)$ and that a maximal compact connected abelian subgroup T of $P \times \{1\}$ is contained in H . Then $H = G$.

Proof. We consider N as in Lemma 2. By Lemma 2(ii) we have $T \subseteq N$. Set $T = T_1 \times \{1\}$ and $N \cap (P \times \{1\}) = N_1$. The maximal compact connected abelian subgroup T_1 of P is contained in N_1 , which by Lemma 2(ii) is a normal subgroup of P . Since all maximal compact connected abelian subgroups are conjugate and their union covers P we conclude that $N_1 = P$. Hence $P \times \{1\} \subseteq N \subseteq H$. Thus H contains the kernel of pr_A . Then from $(*)$ it follows that H is dense in G . As H is closed, $H = G$ follows. ■

Lemma 4. Let P be any compact connected group and T a maximal compact connected abelian subgroup. Let A denote a locally compact group with a topological generating set X . Assume that $X = Y \cup Z \subseteq A$ and $\pi: X \rightarrow P$ are such that $T \cup \pi(Z)$ is a topological generating set of P and that $\{\pi(y), y\} : y \in Y$ topologically generates a subgroup containing $T \times \{1\}$. Then $\text{graph}(\pi) = \{(\pi(x), x) : x \in X\}$ is a topological generating set of $P \times A$.

Proof. Let $H = \overline{\langle X \rangle}$. We must show $H = P \times A$. By Lemma 3 this is the case if the conditions of Lemma 3 hold. Since X is a topological generating set of A then $\text{pr}_A(H)$ is dense in A , and since $\text{pr}_P(H)$ contains T and $\pi(Z)$ it is dense in P . Thus $(*)$ holds. The remaining condition of Lemma 3 is satisfied by hypothesis. ■

We note that every locally compact connected group G contains a characteristic subgroup G_k such that G_k contains the radical R and has no nontrivial compact simple homomorphic image, and is such that G_k/R contains no compact connected normal subgroup. We have $G = G_k$ iff G has no nontrivial compact connected simple group as a homomorphic image. We then say that G is *without compact factors*. Indeed if G is semisimple, then there is a closed semisimple Lie subgroup G_k containing all simple noncompact factors and no compact one. Then G is the product of G_k and the product of all compact simple factors. If G is not semisimple, then G_k is the full inverse image of $(G/R)_k$ in G .

We note that, in particular, if $f: G \rightarrow G^*$ is a quotient homomorphism, then $f(G_k) = G_k^*$.

Proposition 5. *Assume that P is any compact group whose identity component P_0 is contained in $\overline{P'}$, and A is a locally compact connected group. Let X be a topological generating set of A .*

- (i) *If $\pi: X \rightarrow P$ is any function such that $\pi(X)$ is a topological generating set of P then $\text{graph}(\pi) = \{(\pi(x), x) : x \in X\}$ is a topological generating set of a subgroup $H \subseteq P \times A$ containing $\{\mathbf{1}\} \times A_k$.*
- (ii) *If A is without compact factors, then $\text{graph}(\pi)$ topologically generates $P \times A$ for any π as in (i).*

Proof. We let H be the closure of the group generated by $\text{graph}(\pi)$ and denote $P \times A$ by G . We claim $G = H$. Let N be as in Lemma 2. If $H/N = G/N$ then $H = G$. Now $(A/N_A)_k = (A_k N_A)/N_A$ and $(P/N_P)_0 = (P_0 N_P)/N_P$, the latter being contained in $(P/N_P)' = (\overline{P'} N_P)/N_P$. Thus, without loss of generality we can assume $N = \{\mathbf{1}\}$.

We claim that A is isomorphic to P and thus is compact. By Lemma 2, there is a unique injective morphism $f: A \rightarrow P$ extending π with dense image. Thus A is maximally almost periodic and is therefore $V \times C$ with a vector group V and a compact connected group C . Also, P is now a compact, connected group satisfying $P = \overline{P'}$. This last property implies that the identity component of the center of P is trivial. Hence $\text{pr}_P(V)$ is a point and thus V is singleton. Hence $A = C$ is compact, and is mapped isomorphically onto P . Now $A \cong P$ is a compact connected semisimple group and thus $A_k = \{\mathbf{1}\}$. This proves (i).

In order to prove (ii) we note that if $A_k = A$, then after factoring N we have $A = \{\mathbf{1}\}$. After having factored N , we have $P = \{\mathbf{1}\}$. We conclude that $N = G$ and thus $H = G$ by Lemma 2(iii). ■

Certain information obtained in the proof may be of independent interest: If $N = \{\mathbf{1}\}$ then $P \cong A$.

We shall say that a topological group G is (topologically) *perfect* if it satisfies $\overline{G'} = G$. (This holds for example for all locally compact connected groups without radical.)

As a corollary of Proposition 5 we have:

Proposition 6. *Let P be a compact group such that P_0 is perfect. If A is locally compact connected group without compact factors, and if $\sigma(P) \leq \sigma(A)$,*

then $\sigma(P \times A) = \sigma(A)$. ■

We shall now make a few observations about the abelian situation which we shall use presently.

If X is a subset of a locally compact abelian group A , let $F(X)$ denote the free (discrete) abelian group on X and $f: F(X) \rightarrow A$ the morphism extending the inclusion $X \rightarrow A$. The image is dense iff the adjoint morphism $\widehat{f}: \widehat{A} \rightarrow \widehat{F(X)} = \mathbb{T}^X$ is injective. Thus $\sigma(A)$ is the smallest among the cardinals of those sets X for which \widehat{A} has a continuous injective image in \mathbb{T}^X . For instance, if $A = \mathbb{R}^n$, then $\widehat{A} \cong \mathbb{R}^n$, and \mathbb{R}^n has a continuous injective image in \mathbb{T}^{n+1} but not in \mathbb{T}^n . If A is compact and connected then \widehat{A} is discrete and torsion free. It is (algebraically) isomorphic to a subgroup of \mathbb{T}^X if (and only if) the rank of \widehat{A} does not exceed $\mathfrak{c} \cdot \text{card } X$, where \mathfrak{c} is the cardinality of the continuum. Recall that the dimension of A in this case is the rank of \widehat{A} .

Lemma 7. *Assume that K is a compact connected abelian group of dimension $\leq \aleph_0$ and $A = \mathbb{R}^n \times K$ with $n \geq 1$. Let X be a topological generating set of A and assume that C is a compact abelian group with $\sigma(C) \leq \text{card } X$. Then there is a function $\pi: X \rightarrow C$ such that $\{(\pi(x), x) : x \in X\}$ is a topological generating set of $C \times A$.*

Proof. Let $\widehat{f}: \widehat{A} \rightarrow \mathbb{T}^X$ be as in the remarks preceding the lemma. The image $V = \widehat{f}(\mathbb{R}^n \times \{1\})$ has a compact connected closure, and since \widehat{f} is injective the index of V in \overline{V} is of continuum cardinality. The rank of $\widehat{f}(\{0\} \times \widehat{K})$ is countable. Hence its divisible hull K^* is countable and torsion free. Since V is divisible, there is a subgroup W of \mathbb{T}^X such that $\mathbb{T}^X = W \oplus V \oplus K^*$ algebraically. The rank of W is at least \mathfrak{c} , but also it cannot be bigger since $X \subseteq A$ and $\text{card } A = \mathfrak{c}$. Since W contains the torsion group of \mathbb{T}^X and has the same rank as \mathbb{T}^X it is algebraically isomorphic to \mathbb{T}^X under an algebraic isomorphism $i: \mathbb{T}^X \rightarrow W$. Now let $\varphi: (\mathbb{T}^X)_d \oplus \widehat{A} \rightarrow \mathbb{T}^X$ be the injective morphism given by $\varphi(\alpha \oplus \beta) = i(\alpha) + \widehat{f}(\beta)$. Notice that the character group $A(X)$ of $(\mathbb{T}^X)_d$ is the free compact abelian group on the set X . (See [4].) The dual morphism $\widehat{\varphi}: F(X) \rightarrow A(X) \times A$ is given by $\widehat{\varphi}(g) = (\widehat{i}(g), f(g))$ and has dense image. It satisfies $\widehat{\varphi}(x) = (\widehat{i}(x), x)$ where $\{\widehat{i}(x) : x \in X\}$ is topologically generating in $A(X)$. Since $\sigma(C) \leq \text{card}(X)$ there is a surjective homomorphism $\eta: A(X) \rightarrow C$. If we set $\pi = \eta \circ (\widehat{i}|_X)$ then we have the desired function. ■

We obtain at once the following variant of the preceding lemma:

Lemma 7'. *Suppose that K is a compact connected abelian group of dimension $\leq \aleph_0$ and $A = \mathbb{R}^n \times K$ with $n \geq 1$. Let $\{x_j : j \in J\}$ be a topological generating family in A and suppose that C is a compact abelian group with $\sigma(C) \leq \text{card } J$. Then there is a function $\pi: J \rightarrow C$ such that $\{(\pi(j), x_j) : j \in J\}$ is a topological generating set of $C \times A$. ■*

Lemma 8. *If P is a nonsingleton monothetic compact group, then $\sigma(P \times \mathbb{R}^n) = n + 1$.*

Proof. For $n = 0$ the assertion is trivial. If $n \geq 1$ then $\sigma(\mathbb{R}^n) = n + 1$ by the remarks preceding Lemma 7. (There are other arguments for this conclusion: See [3].) Now Lemma 7 implies the conclusion (with $K = \{1\}$). ■

Lemma 9. *Let A denote a locally compact group such that $A/\overline{A'}$ is second countable and connected. Assume that C is a compact abelian group satisfying $\sigma(C) \leq \sigma(A)$. Then for any generating set X of A there is a function $\pi: X \rightarrow P$ such that $\text{graph}(\pi)$ is a generating set of $P \times A$.*

Proof. Let X denote a topological generating set of A and let $f: A \rightarrow A/\overline{A'}$ be the quotient map. Then $\{f(x) : x \in X\}$ is a topological generating family of $f(A)$. By Lemma 7', there is a function $\pi: X \rightarrow C$ such that $\{(\pi(x), f(x)) : x \in X\}$ topologically generates $C \times f(A)$. Let H be the closed subgroup generated by $\{(\pi(x), x) : x \in X\}$ in $C \times A$. We shall prove that $H = G \stackrel{\text{def}}{=} C \times A$ and thereby finish the proof.

Let $h = (\pi(x_1), x_1) \cdots (\pi(x_m), x_m)$ and $h' = (\pi(x'_1), x'_1) \cdots (\pi(x'_n), x'_n)$. Set $[a, b] = aba^{-1}b^{-1}$. Then $[h, h'] = (1, [x_1 \cdots x_m, x'_1 \cdots x'_n])$ since C is abelian. Hence $D = \{1\} \times \overline{A'} \subseteq H$. But if we identify $C \times f(A)$ with G/D , then $H/D = C \times f(A)$ because H/D contains all $(\pi(x), f(x))$, $x \in X$. Then $H = G$ follows. ■

Lemma 10. *Let P be any compact group with identity component $P_0 \subseteq \overline{P'}$ and assume $w(P_0) \leq \mathfrak{c}$. Also assume that A is a locally compact connected group with a second countable factor group A/A_k such that $\sigma(P) \leq \sigma(A)$ and $\sigma(P/P_0) \leq \sigma(A/A_k) - 2$. Then for any topological generating set X of A there is a function $\pi: X \rightarrow P$ such that $\text{graph}(\pi)$ topologically generates $P \times A$.*

Proof. If $A = A_k$ then Proposition 5 proves the assertion. Now assume $A_k \neq A$. We let $f: A \rightarrow A/A_k$ denote the quotient map. If X is a topological generating set of A then $f(X)$ is a topological generating set of A/A_k . Since A/A_k is nonabelian, there are at least two elements y and z such that $f(y) \neq f(z)$.

Let T denote a maximal compact connected abelian subgroup of P . Then there is a $p \in P_0$ such that $T \cup \{p\}$ is a topological generating set of P_0 (see [6], Corollary 2.5). Since $w(P_0) \leq \mathfrak{c}$, then T is monothetic. If M is the closed subgroup generated by $f(y)$ in A/A_k , then there is a $t \in T$ such that $(t, f(y))$ topologically generates $T \times M$ by Lemma 7. Now the closed subgroup topologically generated by $\{(t, f(y)), (p, f(z))\} \cup (\{1\} \times f(X_0))$ with $X_0 = X \setminus \{y, z\}$ has dense projections into A and P_0 and thus generates $P_0 \times A/A_k$ by Lemma 3.

Define $\pi: X \rightarrow P$ by $\pi(y) = t$, $\pi(z) = p$, and such that $\pi(X_0)$ is a generating set of P modulo P_0 . Such a choice is possible because $\sigma(P/P_0) \leq \sigma(A/A_k) - 2 \leq |X_0|$. Let H denote the closed subgroup generated in $G = P \times A$ by $\text{graph}(\pi)$. Let N be defined as in Lemma 2. Then $\{1\} \times A_k$ is contained in N by Proposition 5. Now H/N is topologically generated by $\{(\pi(x), f(x)) : x \in X\} = \{(t, f(y)), (p, f(z))\} \cup \{(\pi(x), f(x)) : x \in X_0\}$ and thus agrees with G/N by the preceding. Now $G = H$ follows. ■

Lemma 11. *Let $\theta: G \rightarrow H$ be an open surjective morphism of locally compact groups such that $\ker \theta$ is compact totally disconnected and contained in G_0 . Then*

- (i) *for each generating set X of H each subset X' of G with $\theta(X') = X$ is a generating subset of G , and*
- (ii) $\sigma(G) = \sigma(H)$.

Proof. We begin by noting that (ii) follows from (i): If X is topological generating set of G with $\text{card } X = \sigma(G)$, then $\theta(X)$ is a topological generating set of H , whence $\sigma(H) \leq \text{card } X = \sigma(G)$. The reverse inclusion, however, we deduce from (i) by taking a generating subset X of H with $\text{card } X = \sigma(H)$ and considering a subset $X' \subseteq G$ such that $\theta|_{X'}: X' \rightarrow X$ is bijective. Then $\sigma(G) \leq \text{card } X' = \text{card } X = \sigma(H)$ by (i).

Now we prove (i). Let X be a topological generating set of H . Let $X' \subseteq G$ be any subset such that $\theta(X') = X$. Let $K = \overline{\langle X' \rangle}$. Then $\theta(K) = H$ since $N \stackrel{\text{def}}{=} \ker \theta$ is compact. It follows that $\theta(K_0) = H_0$. Since N is compact, NK_0 is closed, and since $\theta: G \rightarrow H$ is a proper map, then $\theta|_{NK_0}: NK_0 \rightarrow \theta(K_0)$ is a proper map. Thus $NK_0/N \cong \theta(K_0) = H_0 \cong G_0/N$. We conclude $G_0 = NK_0$. The homogeneous spaces G_0/K_0 and $N/(N \cap N_0)$ are homeomorphic, and the latter is totally disconnected, as N is totally disconnected. The former is connected as G_0 is connected. Thus they are singleton and $K_0 = G_0$. Now let $\Theta: G/G_0 \rightarrow H/H_0$ be defined by $\Theta(gG_0) = \theta(g)H_0$. Then Θ is an isomorphism because $G/G_0 \cong (G/N)/(G_0/N) \cong H/H_0$. Now $\Theta(K/G_0) = \theta(K)/H_0 = H/H_0 = \Theta(G/G_0)$. Since Θ is an isomorphism we conclude $K/G_0 = G/G_0$ and thus $K = G$. Hence X' is a generating set of G .

The principal result

At last we are able to state and prove the main result.

Theorem 12. *Let P be compact group and A a locally compact group satisfying the following hypotheses:*

- (a) *A is connected,*
- (b) $w(A) \leq \aleph_0$ *(that is, A is second countable),*
- (c) $w(P) \leq \mathfrak{c}$,
- (d) $\sigma(P) \leq \sigma(A)$, *and*
- (e) $\sigma(A) \geq 2$ *and* $\sigma(P/P_0) \leq \sigma(A) - 2$.

Then

- (i) *for every generating set X of A there is a function $\pi: X \rightarrow P$ such that $\{(\pi(x), x) : x \in X\}$ is a generating set of $P \times A$, and*
- (ii) $\sigma(P \times A) = \sigma(A)$.

Proof. (i) implies (ii): If we consider X with $\text{card } X = \sigma(A)$ then $\sigma(P \times A) \leq \text{card}(\{(\pi(x), x) : x \in X\}) = \text{card}(X) = \sigma(A)$. Conversely, if Y is a generating set of $P \times A$ with cardinality $\sigma(P \times A)$, then $\text{pr}_A(Y)$ is a generating set of A of cardinality $\leq \text{card}(Y) = \sigma(P \times A)$. Hence $\sigma(A) \leq \sigma(P \times A)$.

Proof of (i):

Case 1: P is abelian. Then the assertion is true by Lemma 9 in view of (a,b,d).

Case 2: The identity component $Z_0(P)$ of the center of P is trivial. Then by Theorem 1.3 in [5] we know that $P_0 = (\overline{P'})_0 \subseteq \overline{P'}$. Then the assertion follows from Lemma 10 in view of (a,b,c,d,e).

Case 3: P is arbitrary. By the Theorem of Lee [10] there is a compact zero dimensional group D of P such that $P = P_0D$. The set $(P_0)'D$ is a compact subgroup B since $(P_0)'$ is a compact characteristic subgroup of P_0 and thus is normal in P . We have $P = Z_0(P)B$ (see [5], Theorem 1.3) and $D \stackrel{\text{def}}{=} Z_0(P) \cap B$ is a compact totally disconnected central subgroup of P contained in P_0 . Now A , as a locally compact connected group, is σ -compact. Hence $P \times A$ is σ -compact. Hence Lemma 11 applies to the quotient map $\theta: (P \times A) \rightarrow \frac{P \times A}{D \times \{1\}}$. Assume that we have a generating set X of A and a function $\pi_0: X \rightarrow P/D$ such that $\text{graph}(\pi_0)$ is a generating set for $(P/D) \times A \cong \frac{P \times A}{D \times \{1\}}$. Now define $\pi: X \rightarrow P$ such that $\pi(x)D = \pi_0(x)$ for all $x \in X$. Then $\text{graph}(\pi) = \{(\pi(x), x) : x \in X\}$ maps onto $\text{graph}(\pi_0)$. By Lemma 11 then $\text{graph}(\pi)$ is a generating set of $P \times A$.

We therefore can replace P by P/D and thus assume that P is the direct product of the subgroups $Z_0(P)$ and B . Thus $G = Z_0(P) \times B \times A$. Let X be a generating set of A . By Case(1) we find a function $\pi_1: X \rightarrow Z_0(P)$ such that $\{(\pi_1(x), 1, x) : x \in X\}$ is a generating set of $Z_0(P) \times \{1\} \times A$. Now we notice that

$$B/B_0 \rightarrow \frac{Z_0(B) \times B \times A}{Z_0(B) \times B_0 \times A} \cong P/P_0.$$

Hence $\sigma(B/B_0) = \sigma(P/P_0)$ and all hypotheses are satisfied which allow us to apply Lemma 10 with $Z_0(P) \times \{1\} \times A$ in place of A and B in place of P . This yields a function $\pi_2: X \rightarrow B$ such that $\{(\pi_1(x), \pi_2(x), x) : x \in X\}$ is a generating set for $Z_0(P) \times B \times A$. If we recall $P = Z_0(P) \times B$ and define $\pi: X \rightarrow P$ by $\pi(x) = (\pi_1(x), \pi_2(x))$ we see that this is the assertion of the theorem. ■

Note that if P is connected then the condition $\sigma(P/P_0) \leq \sigma(A) - 2$ is trivially satisfied and by [6] conditions (c) and (e) imply (d). In particular, for the case that A is a connected Lie group, we obtain:

Corollary 13. *Let P denote a compact connected group of weight $\leq \mathfrak{c}$ and L a nonsingleton connected Lie group. Then*

- (i) *for every generating set X of A there is a function $\pi: X \rightarrow P$ such that $\text{graph}(\pi)$ is a generating set of $P \times A$, and*
- (ii) $\sigma(P \times L) = \sigma(L)$.

Proof. (i) If L is compact abelian, then L and $P \times L$ are monothetic and the assertion is true. Otherwise $\sigma(L) \geq 2$ and Theorem 12 applies. (ii) follows from (i) ■

We note that assumptions of the sort of conditions (d) and (e) are necessary in order for the conclusion to hold. The topological rank of the group $\mathbb{Z}(2)^n \times \mathbb{R}$ is $\min\{2, n\}$ which is large if n is large while $\sigma(R) = 2$.

It is interesting to realize that condition (b) cannot be replaced by the condition $w(A) \leq \mathfrak{c}$ without invalidating the theorem. In order to understand this let us consider the following example:

Let P be any nonsingleton compact connected group with $w(P) \leq \mathfrak{c}$. Let $K = \widehat{\mathbb{R}}_d$ denote the universal solenoid. Set $A = \mathbb{R} \times K$ and observe $w(P) \leq \mathfrak{c} = w(K) = w(A)$ and $\sigma(P) = 1$, $\sigma(A) = 2$. Thus hypotheses (a,c,d,e) of Theorem 12 are satisfied. The prescription of a generating set X of two elements of A is equivalent to giving a morphism of locally compact groups $e: \mathbb{Z}^2 \rightarrow A$ with dense image and that is tantamount to giving an injective morphism $\widehat{e}: \widehat{A} \rightarrow \mathbb{T}^2$. But $\widehat{A} = \mathbb{R} \times \widehat{\mathbb{R}}_d$. Hence we can choose \widehat{e} to be an isomorphism of \widehat{A} onto a full torsion free complement of the torsion group $(\mathbb{Q}/\mathbb{Z})^2$ of \mathbb{T}^2 . The prescription of any $\pi: X \rightarrow P$ is the specification of two elements $p, q \in P$. Let $\alpha: \mathbb{Z}^2 \rightarrow P$ denote the unique morphism sending $(1, 0)$ to p and $(0, 1)$ to q . Define $\delta: \mathbb{Z}^2 \rightarrow P \times A$ by $\delta(t) = (\alpha(t), e(t))$. Now $\{(\pi(x), x) : x \in X\}$ is a generating set of $P \times A$ iff δ has dense image. But that means exactly that

$$\widehat{\delta}: \widehat{P} \times \widehat{A} = \widehat{P} \times \mathbb{R} \times \widehat{\mathbb{R}}_d \rightarrow \mathbb{T}^2, \quad \widehat{\delta}(r, s, t) = \widehat{\alpha}(r) + \widehat{e}(s, t)$$

is injective. But since $\widehat{e}(\mathbb{R} \times \widehat{\mathbb{R}}_d)$ is a complement of the torsion subgroup of \mathbb{T}^2 and $\alpha(\widehat{A})$ would have to be a nonzero torsion free subgroup meeting $\widehat{e}(\mathbb{R} \times \widehat{\mathbb{R}}_d)$ trivially, this is patently impossible. Notice that we can even take $P = \mathbb{T}$. ■

We saw that $\sigma(\mathbb{R}^n) = n + 1$ for $n \geq 1$. So Corollary 13 yields:

Corollary 14. (CLEARY and MORRIS [3]) *Let P be a compact connected group of weight $\leq \mathfrak{c}$. Then $\sigma(P \times \mathbb{R}^n) = n + 1$, if $n \geq 1$.* ■

Iwasawa Pairs

There is a “classical” theorem by IWASAWA [8] which says that a locally compact connected group contains a compact normal subgroup N and a local Lie group U commuting elementwise with N such that $(n, u) \mapsto nu: N \times U \rightarrow NU$ is a homeomorphism onto an identity neighborhood. Thus there is a Lie group L with an identity neighborhood isomorphic to U , and we obtain an injective morphism $\varphi: L \rightarrow G$ such that $(n, g) \mapsto n\varphi(g): N \times L \rightarrow G$ is a surjective homomorphism which, by the Open Mapping Theorem, is also open. Let us formalize these remarks as follows:

Lemma 15. *Let G denote a locally compact connected group. Then there is a compact normal subgroup N and a connected Lie group L and an injective morphism $\varphi: L \rightarrow G$ such that*

- (i) $[N, \varphi(L)] = \{\mathbf{1}\}$,
- (ii) $G = N\varphi(L)$, and
- (iii) *there is an identity neighborhood U in L such that $(n, u) \mapsto n\varphi(u): N \times U \rightarrow N\varphi(U)$ is a homeomorphism onto an identity neighborhood of $\mathbf{1}$ such that $[N, \varphi(U)] = \{\mathbf{1}\}$.*

Proof. (i) and (ii) are consequences of (iii), and (iii) is the IWASAWA’s Local Product Theorem (see [8]). ■

Definition 16. If G is a locally compact connected group, then an *Iwasawa pair* is a pair $(N, \varphi: L \rightarrow G)$ with a morphism $\varphi: L \rightarrow G$ satisfying the conditions of Lemma 15. ■

Lemma 17. Let G denote a locally compact connected group with weight $w(G) \leq \mathfrak{c}$. If $(N, \varphi: L \rightarrow G)$ is an Iwasawa pair, then $\sigma(N_0) \leq 2$.

Proof. From $w(N_0) \leq w(G) \leq \mathfrak{c}$ we conclude $\sigma(N_0) \leq 2$. (See [7], 4.13.) ■

We observe that G/N is a Lie group locally isomorphic to L . Let $p: G \rightarrow G/N$ denote the quotient map. The kernel of the covering $\Phi = p \circ \varphi: L \rightarrow G/N$ is $K = \varphi^{-1}(N \cap \varphi(L))$.

For an abelian group A we set $\text{rank } A = \dim_{\mathbb{Q}}(\mathbb{Q} \otimes_{\mathbb{Z}} A)$ and call this number the *torsion free rank* of A .

Lemma 18. Let $\theta: G \rightarrow H$ be a surjective morphism of connected locally compact groups and write $N = \ker \theta$. Assume that S is a closed subgroup of G .

- (i) If $G = NS$ and N is countable, then $S = G$.
- (ii) If N is finite and $\theta(S)$ is dense, then $S = G$
- (iii) If N is finite, then $\sigma(G) = \sigma(H)$.
- (iv) If θ is a covering homomorphism of connected Lie groups then $\sigma(G) \leq \sigma(H) + \text{rank}(\ker \theta)$.

Proof. (i) If $\bigcup_{n \in N} nS = NS = G$, then the locally compact space G is a countable union of homeomorphic closed subsets nS . Then by the Baire Category Theorem, one of them has inner points. Thus the subgroup S has nonempty interior and so is open. Hence it is also closed. As G is connected, $S = G$ follows.

(ii) If N is finite, then NS is a closed subgroup. Since $\theta(S)$ is dense it follows that NS is dense. Thus $NS = G$.

(iii) Let $Y \subseteq H$ denote a generating set in the sense that $H = \overline{\langle Y \rangle}$ and let $X \subseteq \theta^{-1}(Y)$ be any subset such that $\theta|_X: X \rightarrow Y$ is bijective. Define $S = \overline{\langle X \rangle}$. Then $\theta(S)$ is dense in H . As N is finite, (ii) applies and proves the assertion.

(iv) Now N is finitely generated abelian. Its torsion group M is finite. By (iii) we have $\sigma(G) = \sigma(G/M)$ and the covering morphism $\Theta: G/M \rightarrow H$, $\Theta(gM) = \theta(g)$ has a free kernel of rank $\text{rank } N$ which is also $\sigma(N)$. Since always $\sigma(G/M) \leq \sigma(H) + \sigma(\ker \Theta)$ (cf. [3]), the assertion follows. ■

Conclusion (iii) also follows from Lemma 11.

For the quotient map $f: \mathbb{R} \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$ we have $\sigma(\mathbb{R}) = 2$, $\sigma(\mathbb{T}) = 1$ and $\text{rank}(\mathbb{Z}) = 1$. This shows that the inequality in (iv) is sharp.

For an Iwasawa pair $(N, \varphi: L \rightarrow G)$ we write $\beta(N, \varphi) = \text{rank}(\ker \Phi)$ and call it its *Betti number*.

Proposition 19. *If G is a noncompact locally compact connected group of weight not exceeding the cardinality of the continuum, then for any Iwasawa pair $(N, \varphi: L \rightarrow G)$,*

$$\sigma(G) \leq \sigma(L) \leq \sigma(G/N) + \beta(N, \varphi) \leq \sigma(G) + \beta(N, \varphi).$$

Proof. Since G is not compact then L is not compact, in particular, L is nonsingleton. Then Corollary 13 shows $\sigma(N_0 \times L) = \sigma(L)$. We consider the surjective morphism $\pi: N \times L \rightarrow G$, $\pi(n, g) = n\varphi(g)$. In Lemma 17 of [7] we show that the arc component G_a of the identity in G is $N_a\varphi(L)$ where N_a is the arc component of $\mathbf{1}$ in N . But G_a is dense in G and $G_a \subseteq N_0\varphi(L) = \pi(N_0 \times L)$. Thus π maps $N_0 \times L$ onto a dense subgroup of G . Hence any topological generating set X of $N_0 \times L$ is mapped onto a topological generating set $\pi(X)$ of G . Therefore, $\sigma(G) \leq \text{card } \pi(X) \leq \text{card } X = \sigma(N_0 \times L) = \sigma(L)$.

Notice that we always have $\sigma(G/N) \leq \sigma(G)$.

In order to prove the remaining inequality we consider the quotient map $q: G \rightarrow G/N$ and the morphism $f \stackrel{\text{def}}{=} q \circ \varphi: L \rightarrow G \rightarrow G/N$. Now $\ker f = \{g \in L : \varphi(g) \in N\}$. If $U \subseteq L$ is as in Lemma 15(iii), then $\varphi(U) \cap N = \{1\}$ and thus $U \cap \ker f = \{1\}$. Thus $\ker f$ is discrete. Because of $G = N\varphi(L)$ the morphism f is surjective, and since L is connected, hence σ -compact, it is open. Thus f is a covering homomorphism. Now Lemma 18(iv) applies and shows the assertion. ■

In order to illustrate the situation by an example we define $N = \widehat{\mathbb{Q}/\mathbb{Z}} \cong \prod_{p \text{ prime}} \mathbb{Z}_p$ where \mathbb{Z}_p is the additive group of p -adic integers. Then $q: \mathbb{Q}/\mathbb{Z} \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$ gives an injective morphism $\widehat{q}: \mathbb{Z} \rightarrow N$ with dense image. Write $i: \mathbb{Z} \rightarrow N \times \mathbb{R}$, $i(n) = (\widehat{q}(n), -n)$. Then i is an injective morphism with discrete image (since the projection of the image onto \mathbb{R} is discrete). Define $G = (N \times \mathbb{R})/(\text{im } i)$ and let $p: N \times \mathbb{R} \rightarrow G$ denote the quotient homomorphism. Since the compact space $N \times [0, 1]$ maps onto G under p we know that G is compact. The dual of the exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{i} N \times \mathbb{R} \xrightarrow{p} G \rightarrow 0$$

is the exact sequence

$$0 \rightarrow \widehat{G} \xrightarrow{\widehat{p}} (\widehat{\mathbb{Q}/\mathbb{Z}}) \times \mathbb{R} \xrightarrow{\widehat{i}} \widehat{\mathbb{R}/\mathbb{Z}} \rightarrow 0$$

with $\widehat{i}(q + \mathbb{Z}, r) = q - r + \mathbb{Z}$. Thus $\text{im } \widehat{p} = \ker \widehat{i} = \{(q + \mathbb{Z}, r) : q - r \in \mathbb{Z}\} = \{(q + \mathbb{Z}, q) : q \in \mathbb{Q}\} \cong \mathbb{Q}$. It follows that G is the rational solenoid $\cong \widehat{\mathbb{Q}}$. The image $N^* = p(N \times \{0\})$ in G is an isomorphic copy of N , and $(N^*, \varphi: \mathbb{R} \rightarrow G)$, $\varphi(r) = p(0, r)$ is an Iwasawa pair for G . Since $G = \widehat{\mathbb{Q}}$, then G is monothetic; that is, $\sigma(G) = 1$. We know $\sigma(\mathbb{R}) = 2$. We have $G/N \cong \mathbb{R}/\mathbb{Z}$ and $\ker(\mathbb{R} \rightarrow G \rightarrow G/N) = \{r \in \mathbb{R} : \varphi(r) = p(0, r) \in N^*\} = \{r \in \mathbb{R} : (\exists n \in \mathbb{Z})(0, r) \in (\widehat{q}(n), -n) + (N \times \{0\})\} = \mathbb{Z}$. Thus $\beta(N^*, \varphi) = 1$. Thus the second inequality of Proposition 19 is sharp.

Remarks and Open Questions

There is a result by KURANISHI [9] saying that every connected semisimple Lie group has topological rank 2. This suggests the following question:

Question A. *If G is a perfect connected nondegenerate Lie group is $\sigma(G) = 2$?*

An analysis of the topological rank of a connected Lie group requires an answer to the following question:

Question B. *What is the topological rank of a solvable connected Lie group?*

In this context the following result of CLEARY [1] is relevant:

Proposition 20. *If G is a connected nilpotent Lie group then $\sigma(G) = \sigma(G/\overline{G'})$. ■*

Note that in Proposition 20 $G/\overline{G'} \cong \mathbb{R}^n \times \mathbb{T}^m$ and thus $\sigma(G) = n + 1$.

Finally we mention that in [6,7] we discussed the cardinal invariant $s(G)$ for a locally compact group defined as the minimum cardinal of a suitable subset X of G where X is a topological generating subset of G such that X is discrete and closed in $G \setminus \{1\}$. Trivially, $\sigma(G) \leq s(G)$. Further it was shown in [6] that for a locally compact connected group G of weight $\leq \mathfrak{c}$, the cardinal $s(G)$ is finite and so equals $\sigma(G)$. This leaves open the question whether these two cardinals are always equal. They are not as the example below demonstrates.

Let X be a set of cardinality $2^{\mathfrak{c}}$. Now let $G = A(X)$ be the free compact abelian group on the discrete set X [4]. Then $\widehat{G} = (\mathbb{T}^X)_d$ and thus $w(G) = \text{rank } \widehat{G} = 2^{\mathfrak{c} \cdot 2^{\mathfrak{c}}} = 2^{2^{\mathfrak{c}}}$. In [6] we showed for a compact connected group H with $w(H) > \mathfrak{c}$ that $s(H)^{\aleph_0} = w(H)^{\aleph_0}$. Thus $s(G)^{\aleph_0} = 2^{2^{\mathfrak{c}}}$. On the other hand, X is a topological generating subset of G . Hence $\sigma(G) \leq \text{card } X = 2^{\mathfrak{c}}$. Since $(2^{\mathfrak{c}})^{\aleph_0} = 2^{\mathfrak{c}}$ we conclude that $\sigma(G) < s(G)$.

If $\sigma(G)$ is infinite, then it coincides with the density $d(G)$. The relation $d(G) = \log w(G)$ was proved by COMFORT and ITZKOWITZ in [1].

References

- [1] Comfort, W. W., and G. L. Itzkowitz, *Density characters in topological groups*, Math. An.. **226** (1977), 223–227.
- [cl] Cleary, Joan, “On the structure of locally compact Hausdorff groups”, Ph.D. Thesis, La Trobe University, Australia, 1989, ix+110 pp.
- [3] Cleary, Joan, and Sidney A. Morris, *Generators for locally compact groups*, submitted.
- [4] Hofmann, Karl Heinrich and Sidney A. Morris, *Free compact groups I: Free compact abelian groups*, Topology Appl. **23** (1986), 41–64, and Errata, *ibid* **28** (1988), 101–102.
- [5] —, *Free compact groups II: The center*, Topology Appl. **28** (1988), 215–231.
- [6] —, *Weight and c* , J. Pure Appl. Algebra **68** (1990), 181–194.
- [7] —, *Generators on the arc component of compact connected groups*, submitted.
- [8] Iwasawa, K., *On some types of topological groups*, Ann. of Math. **50** 1949, 507–558.
- [9] Kuranishi, M., *Two element generation on semi-simple Lie groups*, Kōdai Math. Sem. Rep. Nos. 5–6, Dec. 1949.
- [10] Lee, Dong Hoon, *Supplements for the identity component in locally compact groups*, Math. Z. **104** (1968), 28–49.

Fachbereich Mathematik
Technische Hochschule Darmstadt
Schlossgartenstr. 7
D-6100 Darmstadt,
Germany
hofmann@mathematik.th-darmstadt.de

Faculty of Informatics
University of Wollongong
Locked Bag 8844,
South Coast Mail Centre, NSW 2521,
Australia
sid@wampyr.cc.uow.edu.au

Received July 3, 1992